

Discussion Paper Series

RIEB


Kobe University

DP2016-37

**A Generalization of Fatou's Lemma for
Extended Real-Valued Functions on
 σ -Finite Measure Spaces: With an
Application to Infinite-Horizon
Optimization in Discrete Time**

Takashi KAMIHIGASHI

Revised January 10, 2017



Research Institute for Economics and Business Administration

Kobe University

2-1 Rokkodai, Nada, Kobe 657-8501 JAPAN

A Generalization of Fatou's Lemma for Extended Real-Valued Functions on σ -Finite Measure Spaces: With an Application to Infinite-Horizon Optimization in Discrete Time

Takashi Kamihigashi*

January 8, 2017

Abstract

Given a sequence $\{f_n\}_{n \in \mathbb{N}}$ of measurable functions on a σ -finite measure space such that the integral of each f_n as well as that of $\limsup_{n \uparrow \infty} f_n$ exists in $\overline{\mathbb{R}}$, we provide a sufficient condition for the following inequality to hold:

$$\limsup_{n \uparrow \infty} \int f_n d\mu \leq \int \limsup_{n \uparrow \infty} f_n d\mu.$$

Our condition is considerably weaker than sufficient conditions known in the literature such as uniform integrability (in the case of a finite measure) and equi-integrability. As an application, we obtain a new result on the existence of an optimal path for deterministic infinite-horizon optimization problems in discrete time.

Keywords: Fatou's lemma; σ -finite measure space; infinite-horizon optimization; hyperbolic discounting; existence of optimal paths

*RIEB, Kobe University, Rokkodai, Nada, Kobe 657-8501 JAPAN. Email: tkamihig@rieb.kobe-u.ac.jp. Tel/Fax: +81-78-803-7015.

1 Introduction

Let $(\Omega, \mathcal{F}, \mu)$ be a measure space. Let $\mathcal{L}(\Omega)$ be the set of measurable functions $f : \Omega \rightarrow \overline{\mathbb{R}}$. A standard version of (reverse) Fatou's lemma states that given a sequence $\{f_n\}_{n \in \mathbb{N}}$ in $\mathcal{L}(\Omega)$, if there exists an integrable function $f \in \mathcal{L}(\Omega)$ such that $f_n \leq f$ μ -a.e. for all $n \in \mathbb{N}$, then

$$\overline{\lim}_{n \uparrow \infty} \int f_n d\mu \leq \int \overline{\lim}_{n \uparrow \infty} f_n d\mu, \quad (1.1)$$

where $\overline{\lim} = \limsup$. We call the above inequality the *Fatou Inequality*.

Some sufficient conditions for this inequality weaker than the one described above are known. In particular, provided that the integral of each f_n as well as that of $\overline{\lim}_{n \uparrow \infty} f_n$ exists, “uniform integrability” of $\{f_n^+\}$ (where f_n^+ is the positive part of f_n) is a sufficient condition for the Fatou inequality (1.1) in the case of a finite measure (e.g., [6, 4, 7, 17]); so is “equi-integrability” of the same sequence in the case of a σ -finite measure (see [9, 10]). These conditions are precisely defined in Section 2.

In this paper we provide a sufficient condition for the Fatou inequality (1.1) considerably weaker than the above conditions. Our approach is based on the following assumption, which is maintained throughout the paper.

Assumption 1.1. $(\Omega, \mathcal{F}, \mu)$ is a σ -finite measure space.

Under this assumption there is an increasing sequence of measurable sets of finite measure whose union equals Ω . We use this sequence to specify a “direction” in which we successively approximate the integral of a function.

There is a natural increasing sequence of measurable sets if the measure space is the set of nonnegative integers equipped with the counting measure. In this setting, we provide a simple sufficient condition for the Fatou inequality (1.1) as a corollary of our general result. Applying this condition to a fairly general class of infinite-horizon deterministic optimization problems in discrete time, we establish a new result on the existence of an optimal path. The condition takes a form similar to transversality conditions and other related conditions in dynamic optimization (e.g., [11, 12, 13, 14]).

The current line of research was initially motivated by the limitations of the existing applications of Fatou's lemma to dynamic optimization problems (e.g., [5, 8]). In particular, there are certain cases in which optimal paths exist but the standard version of Fatou's lemma fails to apply. This is illustrated with some examples following our existence result.

We should mention that there are other important extensions of Fatou's lemma to more general functions and spaces (e.g., [3, 2, 16]). However, to our knowledge, there is no result in the literature that covers our generalization of Fatou's lemma, which is specific to extended real-valued functions.

In the next section we define the concepts and conditions needed to state our main result and to compare it with some previous results based on uniform integrability and equi-integrability. In Section 3 we state our main result and derive those previous results as consequences. In Section 5 we present two simple examples that cannot be treated by the previous results but that can easily be treated using our result. In Section 6 we show a new result on the existence of an optimal path for infinite-horizon deterministic optimization problems in discrete time. In Section 8 we prove our main result.

2 Definitions

Given $f \in \mathcal{L}(\Omega)$, let f^+ and f^- denote the positive and negative parts of f , respectively; i.e., $f^+ = \max\{f, 0\}$ and $f^- = \max\{-f, 0\}$. A function $f \in \mathcal{L}(\Omega)$ is called *semi-integrable* if f^+ or f^- is integrable, and *upper (lower) semi-integrable* if f^+ (f^-) is integrable.

We say that a sequence $\{A_i\}_{i \in \mathbb{N}}$ in \mathcal{F} is a σ -finite *exhausting sequence* if

$$\forall i \in \mathbb{N}, \quad A_i \subset A_{i+1}, \quad \mu(A_i) < \infty, \quad (2.1)$$

$$\mu(\Omega \setminus \cup_{i \in \mathbb{N}} A_i) = 0. \quad (2.2)$$

It is easy to see that μ is σ -finite if and only if there exists a σ -finite exhausting sequence. Since we assume that μ is σ -finite, we have at least one σ -finite exhausting sequence.

A sequence $\{f_n\}_{n \in \mathbb{N}}$ of integrable functions in $\mathcal{L}(\Omega)$ is called *equi-integrable* (e.g., [10, p. 16]) if the following conditions hold:

- (a) For any $\epsilon > 0$ there exists $\delta > 0$ such that any $A \in \mathcal{F}$ with $\mu(A) < \delta$ satisfies

$$\sup_{n \in \mathbb{N}} \int_A |f_n| d\mu \leq \epsilon. \quad (2.3)$$

- (b) For any $\epsilon > 0$ there exists $E \in \mathcal{F}$ with $\mu(E) < \infty$ such that

$$\sup_{n \in \mathbb{N}} \int_{\Omega \setminus E} |f_n| d\mu \leq \epsilon. \quad (2.4)$$

Suppose that $\mu(\Omega) < \infty$. A sequence $\{f_n\}_{n \in \mathbb{N}}$ of integrable functions in $\mathcal{L}(\Omega)$ is called *uniformly integrable* (e.g., [7, p. 144]) if

$$\lim_{c \uparrow \infty} \left[\sup_{n \in \mathbb{N}} \int_{\{|f_n| \geq c\}} |f_n| d\mu \right] = 0. \quad (2.5)$$

It is known that a sequence $\{f_n\}_{n \in \mathbb{N}}$ of integrable functions in $\mathcal{L}(\Omega)$ is uniformly integrable if and only if $\sup_{n \in \mathbb{N}} \int |f_n| d\mu < \infty$ and condition (a) above holds (e.g., [7, p. 144]). In the case of a finite measure, condition (b) trivially holds, and thus uniform integrability implies equi-integrability. Conversely, provided that $\sup_{n \in \mathbb{N}} \int |f_n| d\mu < \infty$, equi-integrability implies uniform integrability on each measurable set of finite measure; see [10, Proposition 2.8] for related results.

3 A Generalization of Fatou's Lemma

We are ready to state the main result of this paper.

Theorem 3.1. *Let $\{f_n\}_{n \in \mathbb{N}}$ be a sequence of semi-integrable functions in $\mathcal{L}(\Omega)$ such that $\overline{\lim}_{n \uparrow \infty} f_n$ is semi-integrable. Let $\{B_i\}_{i \in \mathbb{N}} \subset \mathcal{F}$ be a σ -finite exhausting sequence. Suppose that*

$$\underline{\lim}_{i \uparrow \infty} \overline{\lim}_{n \uparrow \infty} \int_{\Omega \setminus A_i} f_n d\mu \leq 0 \quad (3.1)$$

for any σ -finite exhausting sequence $\{A_i\}_{i \in \mathbb{N}} \subset \mathcal{F}$ such that

$$(i) \forall i \in \mathbb{N}, A_i \subset B_i, \quad (ii) \lim_{i \uparrow \infty} \mu(B_i \setminus A_i) = 0. \quad (3.2)$$

Then the Fatou inequality (1.1) holds.

Proof. See Section 8. □

It is shown in the proof (Lemma 8.4) that (2.1) and (3.2) imply (2.2); i.e., (2.1) and (3.2) imply that $\{A_i\}_{i \in \mathbb{N}}$ is a σ -finite exhausting sequence. Thus in Theorem 3.1, the requirement that $\{A_i\}$ be a σ -finite exhausting sequence can be replaced with (2.1). However, to verify (3.1) to apply Theorem 3.1, it is useful to have (2.2) instead of deriving it; for example, see the proofs of Corollaries 4.1 and 4.2.

If $\Omega = \mathbb{Z}_+$ and μ is the counting measure, we obtain a simple sufficient condition for the Fatou inequality:

Corollary 3.1. *Suppose that $\Omega = \mathbb{Z}_+$ and that μ is the counting measure. Let $\{f_n\}_{n \in \mathbb{N}}$ be a sequence of semi-integrable functions in $\mathcal{L}(\Omega)$ such that $\overline{\lim}_{n \uparrow \infty} f_n$ is semi-integrable. Suppose further that*

$$\underline{\lim}_{i \uparrow \infty} \overline{\lim}_{n \uparrow \infty} \sum_{t=i}^{\infty} f_n(t) \leq 0, \quad (3.3)$$

where the sum is understood as the Lebesgue integral with respect to the counting measure μ . Then

$$\overline{\lim}_{n \uparrow \infty} \sum_{t=0}^{\infty} f_n(t) \leq \sum_{t=0}^{\infty} \overline{\lim}_{n \uparrow \infty} f_n(t). \quad (3.4)$$

Proof. Assume (3.3). For $i \in \mathbb{N}$, let $B_i = \{0, \dots, i-1\}$. Then $\{B_i\}_{i \in \mathbb{N}}$ is a σ -finite exhausting sequence. Let $\{A_i\}_{i \in \mathbb{N}} \subset \mathcal{F}$ satisfy (3.2). Then $A_i = B_i$ for sufficiently large i . For such i we have

$$\sum_{\Omega \setminus A_i} f_n(t) = \sum_{t=i}^{\infty} f_n(t). \quad (3.5)$$

Hence (3.1) follows from (3.3). Now (3.4) holds by Theorem 3.1. \square

4 Known Extensions of Fatou's Lemma

The version of Fatou's lemma stated at the beginning of this paper can be shown as a consequence of Theorem 3.1.

Corollary 4.1. *Let $\{f_n\}_{n \in \mathbb{N}}$ be a sequence in $\mathcal{L}(\Omega)$ such that for some upper semi-integrable function $f \in \mathcal{L}(\Omega)$ we have $f_n \leq f$ μ -a.e. for all $n \in \mathbb{N}$. Then the Fatou inequality (1.1) holds.*

Proof. Since $f_n \leq f$ μ -a.e. for all $n \in \mathbb{N}$ and f is upper semi-integrable, f_n is upper semi-integrable for each $n \in \mathbb{N}$, and so is $\overline{\lim}_{n \uparrow \infty} f_n$. For any σ -finite exhausting sequence $\{A_i\}_{i \in \mathbb{N}}$ we have

$$\underline{\lim}_{i \uparrow \infty} \overline{\lim}_{n \uparrow \infty} \int_{\Omega \setminus A_i} f_n d\mu \leq \underline{\lim}_{i \uparrow \infty} \int_{\Omega \setminus A_i} f d\mu \leq \underline{\lim}_{i \uparrow \infty} \int_{\Omega \setminus A_i} f^+ d\mu = 0, \quad (4.1)$$

where the equality holds by (2.2) since f is upper semi-integrable. Now the Fatou inequality (1.1) holds by Theorem 3.1. \square

The following version of Fatou's lemma is shown in [6, p. 4] and [4, p. 10], and can be derived as a consequence of Theorem 3.1.

Corollary 4.2. *Suppose that $\mu(\Omega) < \infty$. Let $\{f_n\}_{n \in \mathbb{N}}$ be a sequence of functions in $\mathcal{L}(\Omega)$ such that $\{f_n^+\}_{n \in \mathbb{N}}$ is uniformly integrable. Suppose further that $\overline{\lim}_{n \uparrow \infty} f_n$ is semi-integrable. Then the Fatou inequality (1.1) holds.*

Proof. Recall that uniform integrability of $\{f_n^+\}$ requires integrability of each f_n^+ and condition (a) in Section 2 with f_n^+ replacing f_n . Let $\{A_i\}_{i \in \mathbb{N}}$ be any σ -finite exhausting sequence. We have

$$\liminf_{i \uparrow \infty} \overline{\lim}_{n \in \mathbb{N}} \int_{\Omega \setminus A_i} f_n d\mu \leq \limsup_{i \uparrow \infty} \int_{\Omega \setminus A_i} f_n^+ d\mu = 0, \quad (4.2)$$

where the equality holds by condition (a) since $\{f_n^+\}$ is uniformly integrable and $\lim_{i \uparrow \infty} \mu(\Omega \setminus A_i) = 0$ by (2.2) and the finiteness of μ . Now the Fatou inequality (1.1) holds by Theorem 3.1. \square

The next result is a slight variation on the results shown by [9, Lemma 3.3] and [10, Corollary 3.3]. These latter results (unlike Corollary 4.3 below) do not require upper semi-integrability of $\overline{\lim}_{n \uparrow \infty} f_n$ since they use the upper integral, which always exists, instead of the Lebesgue integral.

Corollary 4.3. *Let $\{f_n\}_{n \in \mathbb{N}}$ be a sequence of integrable functions in $\mathcal{L}(\Omega)$ such that $\{f_n^+\}_{n \in \mathbb{N}}$ is equi-integrable. Suppose that $\overline{\lim}_{n \uparrow \infty} f_n$ is semi-integrable. Then the Fatou inequality (1.1) holds.*

Proof. By equi-integrability of $\{f_n^+\}$ and condition (b) in Section 2, there exists a sequence $\{E_i\}_{i \in \mathbb{N}}$ in \mathcal{F} such that $\mu(E_i) < \infty$ for all $i \in \mathbb{N}$ and

$$\limsup_{i \uparrow \infty} \int_{\Omega \setminus E_i} f_n^+ d\mu = 0. \quad (4.3)$$

Since μ is σ -finite, there exists a σ -finite exhausting sequence $\{C_i\}_{i \in \mathbb{N}}$. For $i \in \mathbb{N}$, let $B_i = (\cup_{j=1}^i E_j) \cup C_i$. Then $\{B_i\}_{i \in \mathbb{N}}$ is also a σ -finite exhausting sequence. Let $\{A_i\}_{i \in \mathbb{N}}$ be a sequence in \mathcal{F} satisfying (3.2).

Fix $i \in \mathbb{N}$ for the moment. For each $n \in \mathbb{N}$ we have

$$\int_{\Omega \setminus A_i} f_n d\mu \leq \int_{\Omega \setminus A_i} f_n^+ d\mu = \int_{\Omega \setminus B_i} f_n^+ d\mu + \int_{B_i \setminus A_i} f_n^+ d\mu. \quad (4.4)$$

Applying $\sup_{n \in \mathbb{N}}$ to the leftmost and rightmost sides, we obtain

$$\sup_{n \in \mathbb{N}} \int_{\Omega \setminus A_i} f_n \leq \sup_{n \in \mathbb{N}} \int_{\Omega \setminus B_i} f_n^+ d\mu + \sup_{n \in \mathbb{N}} \int_{B_i \setminus A_i} f_n^+ d\mu. \quad (4.5)$$

The first supremum on the right-hand side converges to zero as $i \uparrow \infty$ by (4.3) since $E_i \subset B_i$ for all $i \in \mathbb{N}$. The second supremum also converges to zero as $i \uparrow \infty$ by (3.2)(ii) and condition (a) in Section 2. It follows that (3.1) holds for any sequence $\{A_i\}_{i \in \mathbb{N}}$ in \mathcal{F} satisfying (3.2); thus by Theorem 3.1, the Fatou inequality (1.1) holds. \square

5 Examples

In each of the examples below, Ω is taken to be an interval in \mathbb{R} . Accordingly, \mathcal{F} is taken to be the σ -algebra of Lebesgue measurable subsets of Ω , and μ the Lebesgue measure restricted to \mathcal{F} .

Our first example shows that Theorem 3.1 is a strict generalization of Corollaries 4.2 and 4.3 even in the case of a finite measure.

Example 5.1. Let $\Omega = [-1, 1] \setminus \{0\}$. For $n \in \mathbb{N}$, define $f_n : \Omega \rightarrow \mathbb{R}$ by

$$f_n(\omega) = \begin{cases} 0 & \text{if } \omega \in [-1, -1/n), \\ -n & \text{if } \omega \in [-1/n, 0), \\ n & \text{if } \omega \in (0, 1/n], \\ 0 & \text{if } \omega \in (1/n, 1]. \end{cases} \quad (5.1)$$

It is easy to see that there is no upper semi-integrable function that dominates $\{f_n\}_{n \in \mathbb{N}}$; thus Corollary 4.1 does not apply. Furthermore, $\{f_n^+\}$ is not uniformly integrable; indeed, for any $c \geq 0$ we have

$$\sup_{n \in \mathbb{N}} \int_{\{f_n^+ \geq c\}} f_n^+ d\mu = \sup_{n \in \mathbb{N}: n \geq c} n/n = 1. \quad (5.2)$$

Hence Corollary 4.2, which requires uniform integrability of $\{f_n^+\}$, does not apply either. Neither does Corollary 4.3 since equi-integrability implies uniform integrability on a finite measure space provided that $\sup_{n \in \mathbb{N}} \int |f_n| d\mu < \infty$, which is the case here.

By contrast, Theorem 3.1 easily applies. To see this, note that for each $n \in \mathbb{N}$, f_n is integrable, and so is $\lim_{n \uparrow \infty} f_n$. For $i \in \mathbb{N}$, let

$$B_i = [-1, -1/i] \cup (1/i, 1]. \quad (5.3)$$

Then $\{B_i\}_{i \in \mathbb{N}}$ is a σ -finite exhausting sequence. Let $\{A_i\}_{i \in \mathbb{N}}$ be *any* sequence in \mathcal{F} satisfying (3.2)(i). For each fixed $i \in \mathbb{N}$, for any $n \geq i$, we have $f_n = 0$ on B_i , and $\int_{\Omega \setminus A_i} f_n d\mu = \int_{\Omega \setminus B_i} f_n d\mu = 0$. Thus the left-hand side of (3.1) is zero. Hence the Fatou inequality (1.1) holds by Theorem 3.1.

In fact $\int f_n d\mu = 0$ for all $n \in \mathbb{N}$, and $\lim_{n \uparrow \infty} f_n = 0$. Thus both sides of the Fatou inequality (1.1) equal zero.

In the next example, μ is not finite, and the sequence $\{f_n\}_{n \in \mathbb{N}}$ is uniformly bounded from below.

Example 5.2. Let $\Omega = \mathbb{R}_+$. For $n \in \mathbb{N}$, define $f_n : \Omega \rightarrow \mathbb{R}$ by

$$f_n(\omega) = \begin{cases} 0 & \text{if } \omega \in [0, n), \\ n & \text{if } \omega \in [n, n+1), \\ -1 & \text{if } \omega \in [n+1, 2n+1), \\ 0 & \text{if } \omega \geq 2n+1. \end{cases} \quad (5.4)$$

It is easy to see that there is no upper semi-integrable function that dominates $\{f_n\}_{n \in \mathbb{N}}$; thus Corollary 4.1 does not apply.

For any $\delta \in (0, 1)$ we have

$$\int_{[n, n+\delta)} f_n^+ d\mu = n\delta \uparrow \infty \quad \text{as } n \uparrow \infty. \quad (5.5)$$

Thus $\{f_n^+\}$ does not satisfy condition (a) in Section 2. To consider condition (b), let $E \in \mathcal{F}$ with $\mu(E) < \infty$. Then

$$\mu(E) = \sum_{n \in \mathbb{Z}_+} \mu(E \cap [n, n+1)) < \infty, \quad (5.6)$$

which implies that $\lim_{n \uparrow \infty} \mu(E \cap [n, n+1)) = 0$. It follows that

$$\int_{\Omega \setminus E} f_n^+ d\mu = n(1 - \mu(E \cap [n, n+1))) \rightarrow \infty \quad \text{as } n \uparrow \infty. \quad (5.7)$$

Hence $\{f_n^+\}$ does not satisfy condition (b) either. Therefore $\{f_n^+\}$ is far from being equi-integrable; as a consequence, Corollary 4.3 does not apply.

To see that Theorem 3.1 applies, note that for each $n \in \mathbb{N}$, f_n is integrable for each n , and so is $\lim_{n \uparrow \infty} f_n$. For $i \in \mathbb{N}$, let $B_i = [0, i)$. Then $\{B_i\}_{i \in \mathbb{N}}$ is a σ -finite exhausting sequence. Take *any* sequence $\{A_i\}_{i \in \mathbb{N}}$ in \mathcal{F} satisfying (3.2)(i). Then for each fixed $i \in \mathbb{N}$ we have $\int_{\Omega \setminus A_i} f_n d\mu = 0$ for all $n \geq i$. Thus the left-hand side of (3.1) equals zero. Hence the Fatou inequality (1.1) holds by Theorem 3.1.

In fact, as in the previous example, we have $\int f_n d\mu = 0$ for all $n \in \mathbb{N}$, and $\lim_{n \uparrow \infty} f_n = 0$; thus both sides of the Fatou inequality (1.1) equal zero.

6 An Application to Infinite-Horizon Optimization in Discrete Time

In this section we consider a fairly general class of infinite-horizon maximization problems, establishing a new result on the existence of an optimal path using Corollary 3.1. We start with some notation.

For $t \in \mathbb{Z}_+$, let X_t be a metric space. For $t \in \mathbb{Z}_+$, let $\Gamma_t : X_t \rightarrow X_{t+1}$ be a compact-valued upper hemicontinuous correspondence in the sense that for each $x \in X_t$, $\Gamma_t(x)$ is a nonempty compact subset of X_{t+1} , and for any convergent sequence $\{x_n\}_{n \in \mathbb{N}}$ in X_t with limit $x^* \in X_t$ and any sequence $\{y_n\}_{n \in \mathbb{N}}$ with $y_n \in \Gamma_t(x_n)$ for all $n \in \mathbb{N}$, there exists a convergent subsequence $\{y_{n_i}\}_{i \in \mathbb{N}}$ of $\{y_n\}_{n \in \mathbb{N}}$ with limit $y^* \in \Gamma_t(x^*)$; see [18, p. 56] and [1, p. 564] concerning this definition of upper hemicontinuity. For $t \in \mathbb{Z}_+$, let

$$D_t = \{(x, y) \in X_t \times X_{t+1} : y \in \Gamma_t(x)\}. \quad (6.1)$$

For $t \in \mathbb{Z}_+$, let $r_t : D_t \rightarrow \mathbb{R} \cup \{-\infty\}$ be an upper semicontinuous function.

Consider the following maximization problem:

$$\max_{\{x_t\}_{t=1}^{\infty}} \sum_{t=0}^{\infty} r_t(x_t, x_{t+1}) \quad (6.2)$$

$$\text{s.t. } x_{t+1} \in \Gamma_t(x_t), \quad \forall t \in \mathbb{Z}_+, \quad (6.3)$$

$$x_0 \in X_0 \text{ given.} \quad (6.4)$$

We say that a sequence $\{x_t\}_{t=1}^{\infty}$ is a *feasible path* (from x_0) if it satisfies (6.3). We say that a feasible path $\{x_t^*\}_{t=1}^{\infty}$ is *optimal* (from x_0) if for any feasible

path $\{x_t\}_{t=1}^\infty$, we have

$$\sum_{t=0}^{\infty} r_t(x_t, x_{t+1}) \leq \sum_{t=0}^{\infty} r_t(x_t^*, x_{t+1}^*), \quad (6.5)$$

where $x_0^* = x_0$. For the above inequality to make sense, we assume the following.

Assumption 6.1. For each feasible path $\{x_t\}_{t=1}^\infty$, we have

$$\sum_{t=0}^{\infty} \max\{r_t(x_t, x_{t+1}), 0\} < \infty. \quad (6.6)$$

In other words, the mapping $r_t(x_t, x_{t+1}) : t \mapsto \mathbb{R} \cup \{-\infty\}$ is upper semi-integrable.

We are ready to show our existence result.

Proposition 6.1. *Let Assumption 6.1 hold. Suppose that for any sequence $\{\{x_t^n\}_{t=1}^\infty\}_{n \in \mathbb{N}}$ of feasible paths, we have*

$$\lim_{i \uparrow \infty} \overline{\lim}_{n \uparrow \infty} \sum_{t=i}^{\infty} r_t(x_t^n, x_{t+1}^n) \leq 0. \quad (6.7)$$

Then there exists an optimal path.

Proof. Let

$$\nu = \sup \sum_{t=0}^{\infty} r_t(x_t, x_{t+1}), \quad (6.8)$$

where the supremum is taken over all feasible paths $\{x_t\}_{t=1}^\infty$. By the definition of ν , there exists a sequence $\{\{x_t^n\}_{t=1}^\infty\}_{n \in \mathbb{N}}$ of feasible paths such that

$$\lim_{n \uparrow \infty} \sum_{t=0}^{\infty} r_t(x_t^n, x_{t+1}^n) = \nu. \quad (6.9)$$

Since $\Gamma_0(x_0)$ is compact, there exists a convergent subsequence $\{x_1^{n_j}\}_{j \in \mathbb{N}}$ of $\{x_1^n\}_{n \in \mathbb{N}}$ with limit $x_1^* \in \Gamma_0(x_0)$. By the definition of upper hemicontinuity, there exists a convergent subsequence of $\{x_2^{n_j}\}_{j \in \mathbb{N}}$ with limit $x_2^* \in \Gamma_1(x_1^*)$.

Continuing this way and using the diagonal argument, we see that there exists a subsequence of $\{\{x_t^n\}_{t=1}^\infty\}_{n \in \mathbb{N}}$, again denoted by $\{\{x_t^n\}_{t=1}^\infty\}_{n \in \mathbb{N}}$, such that for each $t \in \mathbb{N}$, x_t^n converges to some x_t^* as $n \uparrow \infty$, and for each $t \in \mathbb{Z}_+$, $x_{t+1}^* \in \Gamma_t(x_t^*)$. Hence $\{x_t^*\}_{t=1}^\infty$ is a feasible path, which implies that

$$\sum_{t=0}^{\infty} r_t(x_t^*, x_{t+1}^*) \leq \nu. \quad (6.10)$$

To apply Corollary 3.1, let $f_n(t) = r_t(x_t^n, x_{t+1}^n)$ for $t \in \mathbb{Z}_+$. By Assumption 6.1, for each $n \in \mathbb{N}$, $f_n(t)$ is an upper semi-integrable function of $t \in \mathbb{Z}_+$. For $t \in \mathbb{Z}_+$, let $f^*(t) = r_t(x_t^*, x_{t+1}^*)$. Since $\{x_t^*\}_{t=1}^\infty$ is feasible as shown above, $f^*(t)$ is also an upper semi-integrable function of $t \in \mathbb{Z}_+$ by Assumption 6.1. For each $t \in \mathbb{Z}_+$, by upper semicontinuity of r_t we have

$$\overline{\lim}_{n \uparrow \infty} f_n(t) = \overline{\lim}_{n \uparrow \infty} r_t(x_t^n, x_{t+1}^n) \leq r_t(x_t^*, x_{t+1}^*) = f^*(t). \quad (6.11)$$

Since the rightmost side is an upper semi-integrable function of $t \in \mathbb{Z}_+$, so is the leftmost side. Note that (3.3) directly follows from (6.7). Thus we can apply Corollary 3.1 to obtain (3.4), which is written here as

$$\overline{\lim}_{n \uparrow \infty} \sum_{t=0}^{\infty} r_t(x_t^n, x_{t+1}^n) \leq \sum_{t=0}^{\infty} \overline{\lim}_{n \uparrow \infty} r_t(x_t^n, x_{t+1}^n). \quad (6.12)$$

We are ready to show that $\{x_t^*\}_{t=1}^\infty$ is an optimal path. Recall from (6.9) that

$$\nu = \lim_{n \uparrow \infty} \sum_{t=0}^{\infty} r_t(x_t^n, x_{t+1}^n) \quad (6.13)$$

$$\leq \sum_{t=0}^{\infty} \overline{\lim}_{n \uparrow \infty} r_t(x_t^n, x_{t+1}^n) \quad (6.14)$$

$$\leq \sum_{t=0}^{\infty} r_t(x_t^*, x_{t+1}^*), \quad (6.15)$$

where (6.14) uses (6.12), and (6.15) uses (6.11). It follows from (6.13)–(6.15) and (6.10) that $\{x_t^*\}_{t=1}^\infty$ is an optimal path. \square

As a simple consequence of Proposition 6.1, we obtain a result that can be viewed as an abstract version of the existence result shown in [8, Proposition 4.1]; see [15, Theorem 1] for a similar result that requires stronger assumptions.

Corollary 6.1. *Suppose that there exists an integrable function $\bar{f} : \mathbb{Z}_+ \rightarrow \mathbb{R}_+$ such that for any feasible path $\{x_t\}_{t=1}^\infty$, we have*

$$\forall t \in \mathbb{Z}_+, \quad r_t(x_t, x_{t+1}) \leq \bar{f}(t). \quad (6.16)$$

Then there exists an optimal path.

Proof. Note that (6.16) implies Assumption 6.1. Thus to apply Proposition 6.1, it suffices to verify (6.7) for an arbitrary sequence $\{\{x_t^n\}_{t=1}^\infty\}_{n \in \mathbb{N}}$ of feasible paths. Let $\{\{x_t^n\}_{t=1}^\infty\}_{n \in \mathbb{N}}$ be a sequence of feasible paths. Then by (6.16) we have

$$\lim_{i \uparrow \infty} \overline{\lim}_{n \uparrow \infty} \sum_{t=i}^{\infty} r_t(x_t^n, x_{t+1}^n) \leq \lim_{i \uparrow \infty} \overline{\lim}_{n \uparrow \infty} \sum_{t=i}^{\infty} \bar{f}(t) = \lim_{i \uparrow \infty} \sum_{t=i}^{\infty} \bar{f}(t) = 0, \quad (6.17)$$

where the last equality holds by integrability of \bar{f} . It follows that (6.7) holds; hence an optimal path exists by Proposition 6.1. \square

Corollary 6.1 can be shown directly by using Fatou's lemma to conclude (6.12) from (6.16) in the proof of Proposition 6.1. As illustrated in the next section, Proposition 6.1 covers some important cases to which Corollary 6.1 fails to apply.

7 Examples of Optimization Problems

To illustrate the significance of our existence result, we consider two special cases of the following example.

Example 7.1. Let $u : \mathbb{R}_+ \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing, upper semicontinuous function. Let $\delta : \mathbb{R}_+ \rightarrow \mathbb{R}_{++}$ be a strictly decreasing function. Consider the following maximization problem:

$$\max_{\{x_t\}_{t=1}^\infty} \sum_{t=0}^{\infty} \delta(t) u(c_t) \quad (7.1)$$

$$\text{s.t. } c_t + x_{t+1} = x_t, \quad c_t, x_{t+1} \geq 0, \quad \forall t \in \mathbb{Z}_+, \quad (7.2)$$

$$x_0 \in \mathbb{R}_+ \text{ given.} \quad (7.3)$$

In economics, u and δ are known as a utility function and a discount function, respectively. The above maximization problem is a special case of (6.2)–(6.4) such that for all $t \in \mathbb{Z}_+$, $X_t = \mathbb{R}_+$ and

$$r_t(x, y) = \delta(t)u(x - y), \quad (7.4)$$

$$\Gamma_t(x) = \{y \in \mathbb{R}_+ : 0 \leq y \leq x\}. \quad (7.5)$$

It is easy to see from (7.2) that

$$\forall t \in \mathbb{Z}_+, \quad c_t, x_t \leq x_0. \quad (7.6)$$

For simplicity, we assume that there exists $\theta > 0$ such that

$$(i) \forall c \geq 0, \quad u(c) \leq \theta c, \quad (ii) \quad u(x_0) > 0. \quad (7.7)$$

(Condition (ii) above does not depend on θ .) It is easy to see that condition (i) above implies Assumption 6.1; see (7.13)–(7.16) for details.

Example 7.2. Consider Example 7.1. Most discrete-time economic models assume an exponential discount function of the form

$$\forall t \in \mathbb{Z}_+, \quad \delta(t) = \beta^t \quad (7.8)$$

for some $\beta \in (0, 1)$. In this case, Corollary 6.1 easily applies. To see this, let $\bar{f}(t) = \beta^t u(x_0)$ for $t \in \mathbb{Z}_+$. Then $\bar{f} : \mathbb{Z}_+ \rightarrow \mathbb{R}_+$ is integrable, and (6.16) holds by (7.6). Hence an optimal path exists by Corollary 6.1.

Example 7.3. Consider Example 7.1 again. Although exponential discounting (7.8) is technically convenient (implying time consistency), experimental evidence suggests that “hyperbolic discounting” is more plausible; see, e.g., [19, p. 1]. A simple hyperbolic discount function can be specified as follows:

$$\forall t \in \mathbb{Z}_+, \quad \delta(t) = \frac{1}{1 + \alpha t} \quad (7.9)$$

for some $\alpha > 0$.

In this example, Corollary 6.1 does not apply since there exists no integrable function $\bar{f} : \mathbb{Z}_+ \rightarrow \mathbb{R}_+$ satisfying (6.16) for all feasible paths. To see this, define the feasible path $\{\tilde{x}_t^n\}_{t=1}^\infty$ for each $n \in \mathbb{N}$ by

$$\tilde{x}_t^n = \begin{cases} x_0 & \text{if } t \leq n, \\ 0 & \text{if } t \geq n + 1. \end{cases} \quad (7.10)$$

Then

$$r_t(\tilde{x}_t^n, \tilde{x}_{t+1}^n) = \begin{cases} u(x_0)/(1 + \alpha t) & \text{if } t = n, \\ u(0)/(1 + \alpha t) & \text{otherwise.} \end{cases} \quad (7.11)$$

Hence any \bar{f} satisfying (6.16) must satisfy

$$\bar{f}(t) \geq u(x_0)/(1 + \alpha t), \quad \forall t \in \mathbb{Z}_+. \quad (7.12)$$

Since the right-hand side is not upper semi-integrable in $t \in \mathbb{Z}_+$ by (7.7)(ii), there exists no integrable function \bar{f} satisfying (6.16) for all feasible paths. Hence Corollary 6.1 does not apply.

However, Proposition 6.1 still applies. To see this, let $\{\{x_t^n\}_{t=1}^\infty\}_{n \in \mathbb{N}}$ be a sequence of feasible paths. For any $n, i \in \mathbb{N}$ we have

$$\sum_{t=i}^{\infty} r_t(x_t^n, x_{t+1}^n) = \sum_{t=i}^{\infty} \frac{u(x_t^n - x_{t+1}^n)}{1 + \alpha t} \quad (7.13)$$

$$\leq \sum_{t=i}^{\infty} \frac{\theta(x_t^n - x_{t+1}^n)}{1 + \alpha i} \quad (7.14)$$

$$= \frac{\theta \sum_{t=i}^{\infty} (x_t^n - x_{t+1}^n)}{1 + \alpha i} \quad (7.15)$$

$$\leq \frac{\theta x_i^n}{1 + \alpha i} \leq \frac{\theta x_0}{1 + \alpha i}, \quad (7.16)$$

where (7.14) uses (7.7)(i), and the second inequality in (7.16) uses (7.6). It follows that

$$\lim_{i \uparrow \infty} \overline{\lim}_{n \uparrow \infty} \sum_{t=i}^{\infty} r_t(x_t^n, x_{t+1}^n) \leq \lim_{i \uparrow \infty} \frac{\theta x_0}{1 + \alpha i} = 0. \quad (7.17)$$

Thus (6.7) holds; hence an optimal path exists by Proposition 6.1.

In the above example, the hyperbolic discount function (7.9) is used to show that Corollary 6.1 does not apply. The only property of the discount function required to apply Proposition 6.1 is the equality in (7.17). We summarize this observation in the following example.

Example 7.4. Consider Example 7.1 again. Suppose that

$$\lim_{t \uparrow \infty} \delta(t) = 0. \quad (7.18)$$

Then the argument of Example 7.3 shows that an optimal path exists by Proposition 6.1.

8 Proof of Theorem 3.1

8.1 Preliminaries

Throughout the proof, we fix $\{f_n\}_{n \in \mathbb{N}}$ and $\{B_i\}_{i \in \mathbb{N}}$ to be given by Theorem 3.1. Define $f^* = \overline{\lim}_{n \uparrow \infty} f_n$. For $n \in \mathbb{N}$, define $\hat{f}_n = \sup_{m \geq n} f_m$. We have

$$f^* = \lim_{n \uparrow \infty} \hat{f}_n. \quad (8.1)$$

The following observation helps simplify the proof.

Lemma 8.1. *If f^* is not upper semi-integrable, then the Fatou inequality (1.1) holds.*

Proof. Suppose that f^* is not upper semi-integrable. Then $\int (f^*)^+ d\mu = \infty$, and f^* must be lower semi-integrable (i.e., $\int (f^*)^- d\mu < \infty$) since f^* is semi-integrable by hypothesis. It follows that $\int f^* d\mu = \int (f^*)^+ d\mu - \int (f^*)^- d\mu = \infty$. Thus the Fatou inequality (1.1) trivially holds. \square

Since the above result covers the case in which f^* is not upper semi-integrable, we assume the following for the rest of the proof.

Assumption 8.1. f^* is upper semi-integrable.

8.2 Lemmas

We establish three lemmas before completing the proof of Theorem 3.1.

Lemma 8.2. *Suppose that there exists a σ -finite exhausting sequence $\{A_i\}_{i \in \mathbb{N}}$ satisfying (3.1) and the following:*

$$\forall i \in \mathbb{N}, \quad \overline{\lim}_{n \uparrow \infty} \int_{A_i} f_n d\mu \leq \int_{A_i} f^* d\mu. \quad (8.2)$$

Then the Fatou inequality (1.1) holds.

Proof. Since each f_n is semi-integrable, we have

$$\forall i, n \in \mathbb{N}, \quad \int f_n d\mu = \int_{A_i} f_n d\mu + \int_{\Omega \setminus A_i} f_n d\mu. \quad (8.3)$$

By (3.1) there exists a subsequence $\{A_{i_k}\}_{k \in \mathbb{N}}$ of $\{A_i\}_{i \in \mathbb{N}}$ such that

$$\forall k \in \mathbb{N}, \quad \overline{\lim}_{n \uparrow \infty} \int_{\Omega \setminus A_{i_k}} f_n d\mu < \infty, \quad (8.4)$$

$$\lim_{k \uparrow \infty} \overline{\lim}_{n \uparrow \infty} \int_{\Omega \setminus A_{i_k}} f_n d\mu \leq 0. \quad (8.5)$$

Note that $\{A_{i_k}\}_{k \in \mathbb{N}}$ is a σ -finite exhausting sequence.

Fix $k \in \mathbb{N}$ for the moment. Replacing i with i_k in (8.3) and applying $\overline{\lim}_{n \uparrow \infty}$ to both sides of the resulting equation, we obtain

$$\overline{\lim}_{n \uparrow \infty} \int f_n d\mu = \overline{\lim}_{n \uparrow \infty} \left[\int_{A_{i_k}} f_n d\mu + \int_{\Omega \setminus A_{i_k}} f_n d\mu \right] \quad (8.6)$$

$$\leq \overline{\lim}_{n \uparrow \infty} \int_{A_{i_k}} f_n d\mu + \overline{\lim}_{n \uparrow \infty} \int_{\Omega \setminus A_{i_k}} f_n d\mu \quad (8.7)$$

$$\leq \int_{A_{i_k}} f^* d\mu + \overline{\lim}_{n \uparrow \infty} \int_{\Omega \setminus A_{i_k}} f_n d\mu, \quad (8.8)$$

where (8.7) holds by (8.4), and (8.8) uses (8.2).

Since f^* is upper semi-integrable and $\{A_{i_k}\}_{k \in \mathbb{N}}$ is a σ -finite exhausting sequence, we have $\lim_{k \uparrow \infty} \int_{A_{i_k}} f^* d\mu = \int f^* d\mu < \infty$. Thus applying $\lim_{k \uparrow \infty}$ to the right-hand side of (8.8) yields

$$\overline{\lim}_{n \uparrow \infty} \int f_n d\mu \leq \int f^* d\mu + \lim_{k \uparrow \infty} \overline{\lim}_{n \uparrow \infty} \int_{\Omega \setminus A_{i_k}} f_n d\mu \leq \int f^* d\mu, \quad (8.9)$$

where the last inequality uses (8.5). The Fatou inequality (1.1) follows. \square

Lemma 8.3. *Let $\{A_i\}_{i \in \mathbb{N}}$ be a sequence in \mathcal{F} such that for each $i \in \mathbb{N}$, $\mu(A_i) < \infty$ and \hat{f}_n^+ converges to $(f^*)^+$ uniformly on A_i as $n \uparrow \infty$. Then $\{A_i\}_{i \in \mathbb{N}}$ satisfies (8.2).*

Proof. Let $i \in \mathbb{N}$. Let $\delta > 0$. Since \hat{f}_n^+ converges to $(f^*)^+$ uniformly on A_i as $n \uparrow \infty$, for sufficiently large $n \in \mathbb{N}$ we have $f_n \leq \hat{f}_n \leq \hat{f}_n^+ \leq (f^*)^+ + \delta$ on A_i . Since $(f^*)^+$ is integrable by Assumption 8.1 and $\mu(A_i) < \infty$, (8.2) holds by Fatou's lemma. \square

Lemma 8.4. *Let $\{A_i\}_{i \in \mathbb{N}}$ be a sequence in \mathcal{F} satisfying (2.1) and (3.2). Then $\{A_i\}$ is a σ -finite exhausting sequence.*

Proof. Since $\{A_i\}$ satisfies (2.1) by hypothesis, it suffices to verify (2.2). For any $i, j \in \mathbb{N}$ with $i \leq j$, by (2.1) for $\{B_i\}$, we have

$$\mu(B_i \setminus A_j) \leq \mu(B_j \setminus A_j) \rightarrow 0 \quad \text{as } j \uparrow \infty, \quad (8.10)$$

where the convergence holds by (3.2). It follows that

$$\forall i \in \mathbb{N}, \quad \mu(B_i \setminus \cup_{j \in \mathbb{N}} A_j) = \lim_{j \uparrow \infty} \mu(B_i \setminus A_j) = 0. \quad (8.11)$$

Therefore

$$\mu(\cup_{i \in \mathbb{N}} B_i \setminus \cup_{j \in \mathbb{N}} A_j) = \lim_{i \uparrow \infty} \mu(B_i \setminus \cup_{j \in \mathbb{N}} A_j) = 0. \quad (8.12)$$

Since $\cup_{i \in \mathbb{N}} A_i \subset \cup_{i \in \mathbb{N}} B_i$, we have

$$\mu(\Omega \setminus \cup_{i \in \mathbb{N}} A_i) = \mu(\Omega \setminus \cup_{i \in \mathbb{N}} B_i) + \mu(\cup_{i \in \mathbb{N}} B_i \setminus \cup_{i \in \mathbb{N}} A_i) = 0, \quad (8.13)$$

where the last equality holds by (2.2) for $\{B_i\}$ and (8.12). It follows that $\{A_i\}$ satisfies (2.2). \square

8.3 Completing the Proof of Theorem 3.1

Note from (8.1) that $(f^*)^+ = \lim_{n \uparrow \infty} \hat{f}_n^+$. Let $\{\epsilon_i\}_{i \in \mathbb{N}}$ be a sequence in \mathbb{R}_{++} such that $\lim_{i \uparrow \infty} \epsilon_i = 0$. For each $i \in \mathbb{N}$, by Egorov's theorem there exists $E_i \in \mathcal{F}$ such that $E_i \subset B_i$, $\mu(B_i \setminus E_i) < \epsilon_i$, and \hat{f}_n^+ converges to $(f^*)^+$ uniformly on E_i as $n \uparrow \infty$. For $i \in \mathbb{N}$, let

$$A_i = \cup_{j=1}^i E_j \subset B_i. \quad (8.14)$$

Then for each $i \in \mathbb{N}$, \hat{f}_n^+ converges to $(f^*)^+$ uniformly on A_i as $n \uparrow \infty$. Thus (8.2) holds by Lemma 8.3.

Note that $\{A_i\}_{i \in \mathbb{N}}$ satisfies (2.1) and (3.2) by construction. Thus by Lemma 8.4, $\{A_i\}$ is a σ -finite exhausting sequence. Hence (3.1) holds by the hypothesis of Theorem 3.1. Since (8.2) also holds as shown in the previous paragraph, the Fatou inequality (1.1) holds by Lemma 8.2.

9 Conclusions

In this paper we have provided a sufficient condition for what we call the Fatou inequality:

$$\overline{\lim}_{n \uparrow \infty} \int f_n d\mu \leq \int \overline{\lim}_{n \uparrow \infty} f_n d\mu.$$

Our condition is considerably weaker than sufficient conditions known in the literature such as uniform integrability (in the case of a finite measure) and equi-integrability. We have illustrated the strength of our condition with simple examples. As an application, we have shown a new result on the existence of an optimal path for deterministic infinite-horizon optimization problems in discrete time. We have illustrated the strength of this existence result with concrete examples of optimization problems.

10 Competing Interests

The author declares that he has no competing interests.

11 Acknowledgements

Financial support from the Japan Society for the Promotion of Science (JSPS KAKENHI Grant Number 15H05729) is gratefully acknowledged.

References

- [1] Aliprantis, C.D., Border, K.C., 2006, *Infinite Dimensional Analysis: A Hitchhiker's Guide*, 3rd ed., Springer-Verlag, Berlin.
- [2] Balder, E.J., 1984, A unifying note on Fatou's lemma in several dimensions, *Mathematics of Operations Research* 9, 267–275.
- [3] Brezis, H., Lieb, E., 1983, A relation between pointwise convergence of functions and convergence of functionals, *Proceedings of the American Mathematical Society* 88, 486–490.
- [4] Cairolì, R., Dalang, R.C., 1996, *Sequential Stochastic Optimization*, John Wiley & Sons, New York.

- [5] Cesari, L., 1983, Optimization–Theory and Applications, Springer-Verlag, New York.
- [6] Chow, Y.S., Robbins, H., Siegmund, D., 1971, Great Expectations: The Theory of Optimal Stopping, Houghton Mifflin Company, Boston.
- [7] Denkowski, Z., Migórski, S., Papageorgiou, N.S., 2003, An Introduction to Nonlinear Analysis: Theory, Springer Science+Business Media, New York.
- [8] Ekeland, I., Scheinkman, J.A., 1986, Transversality conditions for some infinite horizon discrete time optimization problems, *Mathematics of Operations Research* 11, 216–229.
- [9] Giner, E., 2009, Calmness properties and contingent subgradients of integral functions on Lebesgue spaces L_p , $1 \leq p < \infty$, *Set-Valued Analysis* 17, 223–243.
- [10] Giner, E., 2012, Fatou’s lemma and lower epi-limits of integral functions, *Journal of Mathematical Analysis and Applications* 394, 13–29.
- [11] Kamihigashi, T., 2001, Necessity of transversality conditions for infinite horizon problems, *Econometrica* 69, 995–1012.
- [12] Kamihigashi, T., 2003, Necessity of transversality conditions for stochastic problems, *Journal of Economic Theory* 109, 140–149.
- [13] Kamihigashi, T., 2005, Necessity of transversality conditions for stochastic models with bounded or CRRA utility, *Journal of Economic Dynamics and Control* 29, 1313–1329.
- [14] Kamihigashi, T., 2014, Elementary results on solutions to the Bellman equation of dynamic programming: existence, uniqueness, and convergence, *Economic Theory* 56, 251–273.
- [15] Le Van, C., Morhaim, L., 2002, Optimal growth models with bounded or unbounded returns: a unifying approach, *Journal of Economic Theory* 105, 158–187.
- [16] Loeb, P.A., Sun, Y., 2007, A general Fatou lemma, *Advances in Mathematics* 213, 741–762.

- [17] Shiryaev, A.N., 2008, *Optimal Stopping Rules*, Springer-Verlag, Berlin.
- [18] Stokey, N., Lucas, R.E., Jr., 1989, *Recursive Methods in Economic Dynamics*, Harvard University Press, Cambridge MA.
- [19] Zarr, N., Alexander, W.H., Brown, J.W., 2014, Discounting of reward sequences: a test of competing formal models of hyperbolic discounting, *Frontiers in Psychology* 5, 1–9.