

Discussion Paper Series

RIEB

Kobe University

DP2013-02

Stochastic Stability in Monotone Economies

Takashi KAMIHIGASHI

John STACHURSKI

January 30, 2013



Research Institute for Economics and Business Administration

Kobe University

2-1 Rokkodai, Nada, Kobe 657-8501 JAPAN

Stochastic Stability in Monotone Economies*

Takashi Kamihigashi^a and John Stachurski^b

^a*Research Institute for Economics and Business Administration, Kobe University*

^b*Research School of Economics, Australian National University*

January 30, 2013

Abstract

This paper extends a family of well-known stability theorems for monotone economies to a significantly larger class of models. We provide a set of general conditions for existence, uniqueness and stability of stationary distributions when monotonicity holds. The conditions in our main result are both necessary and sufficient for global stability of monotone economies that satisfy a weak mixing condition introduced in the paper. Through our analysis we develop new insights into the nature and causes of stability and instability.

JEL Classification: C62, C63

Keywords: Stability, monotonicity, stationary equilibria

1 Introduction

The stability results for monotone economies developed in Hopenhayn and Prescott (1992, theorem 2) have become a standard tool for analysis of dynamics and stationary equilibria. For example, Huggett (1993) used their results to study asset distributions in incomplete-market economies with infinitely-lived agents. The same

*We thank Yiyong Cai, Andrew McLennan, Kazuo Nishimura, Kevin Reffett, Kenji Sato, and Cuong Le Van for many helpful comments. The paper was also improved by suggestions from our editor Gadi Barlevy and three anonymous referees. Financial support from the Japan Society for the Promotion of Science and Australian Research Council Discovery Grant DP120100321 is gratefully acknowledged. *Email addresses:* tkamihig@rieb.kobe-u.ac.jp, john.stachurski@anu.edu.au

results were applied to variants of Huggett's model with features such as habit formation, endogenous labor supply, capital accumulation and international trade (Díaz *et al.*, 2003; Joseph and Weitzman, 2003; Pijoan-Mas, 2006; Marcet *et al.*, 2007; Portes, 2009). They were used to study the classical one-sector optimal growth model by Hopenhayn and Prescott (1992), a stochastic endogenous growth model by de Heer (1999), and a small open economy by Chatterjee and Shuklayev (2010). They have been used in a wide range of OLG models with features such as credit rationing (Aghion and Bolton, 1997; Piketty, 1997), human capital (Owen and Weil, 1998; Lloyd-Ellis, 2000; Cardak, 2004; Couch and Morand, 2005; Hidalgo-Cabrillana, 2009), international trade (Ranjan, 2001; Das, 2006), nonconcave production (Morand and Reffett, 2007), and occupational choice (Lloyd-Ellis and Bernhardt, 2000; Antunes and Cavalcanti, 2007). Other well-known applications include variants of Hopenhayn and Rogerson's (1993) model of job turnover (Cabrales and Hopenhayn, 1997; Samaniego, 2008) and Hopenhayn's (1993) model of entry and exit (Cooley and Quadrini, 2001; Samaniego, 2006).¹

While Hopenhayn and Prescott's stability results have proved to be a useful set of tools, there are important economic models to which they do not apply. One problem is that they assume compactness of the state space, a condition that fails to hold in many applications. A typical example is macroeconomic models where exogenous productivity follows an AR(1) process with unbounded shocks. A restriction to compact state spaces also causes difficulties if, for example, we are studying models of the wealth (or income or firm size) distribution, and our interest centers on whether the stationary distribution follows a power law; or if we wish to analyze dynamics of asset prices in a model where tail events can have large impacts on portfolio returns. Furthermore, since compactness of the state space usually requires that the shocks perturbing the state variables must themselves be bounded, it also precludes the use of some standard probability distributions that are routinely used in applications, such as the normal, lognormal, exponential, Pareto, Cauchy, gamma and t -distributions. In summary, a restriction to compact state spaces forces researchers to make modeling assumptions for technical rather than economic or empirical reasons, and impinges on their ability to address important economic

¹It should be noted here that Hopenhayn and Prescott's results were preceded by similar results in Bhattacharya and Lee (1988). Hopenhayn and Prescott's results were obtained independently, rely on different techniques, and provide a useful separate treatment of existence. On the other hand, Bhattacharya and Lee use a more general notion of mixing, and show exponential convergence rates in their stability results. More details are given in the literature review.

questions.

In this paper we show that it is possible to significantly weaken the conditions of earlier monotone stability results. We begin by introducing a mixing condition called “order reversing” that is weaker than the monotone mixing condition used by Hopenhayn and Prescott. We also relax the restriction that the state space be compact and order bounded. In this setting, we obtain general conditions for monotone, order reversing processes to attain global stability. The conditions are also necessary, and hence we are able to fully characterize global stability for monotone economies that satisfy this very weak mixing condition.

Our discussion of mixing extends a long line of earlier results, as the general concept of mixing plays a key role in the theory of stability of stochastic systems. In essence, mixing refers to movement of the state variable through “most” parts of the state space. For example, irreducibility of finite Markov chains is a classical mixing concept, the definition of which is that any point in the state space can eventually be visited from any other point. Models with a low degree of mixing can become trapped in certain regions of the state space. In such a setting, initial conditions can have permanent effects. In terms of stationary outcomes, the permanent effect of initial conditions can lead to multiple stationary distributions in distinct “absorbing” subsets of the state space. Such outcomes violate the definition of global stability.

On the other hand, when mixing is strong, the state travels widely through the state space regardless of where it starts, and, as a result, the effects of initial conditions tend to die out. Because mixing reduces the importance of initial conditions, it tends to make initial differences smaller over time. On a mathematical level, smaller differences can be translated into smaller *distances* in some appropriate metric. For this reason, mixing properties tend to be related to contraction mapping arguments (because contractions are operators that map distinct points closer together). In fact, at least for the Euclidean case, the existence, uniqueness and stability results of Hopenhayn and Prescott (1992) can all be obtained simultaneously via Banach’s contraction mapping theorem (cf., Bhattacharya and Lee, 1988).

These results are simple, elegant and powerful, and, for infinite state spaces, the monotone mixing conditions are often easier to check and more likely to be satisfied than classical irreducibility conditions. At the same time, there is a sense in which the strength of these results is also their weakness: Strong results usually require strong assumptions, and this case is no exception. In particular, the uniform contraction rate present in the Banach contraction theorem requires that some minimal positive rate of mixing occurs from any point in the state space. This works well in

compact state spaces, where the minimum is usually attained at the extremities of the state space. But when the state space is not compact the same approach tends to break down.

In this paper, we develop contraction-type arguments driven by our weak mixing assumption, but without requiring the uniformity of the previous results. Without uniformity, Banach's theorem does not apply, so we develop a new fixed point result that gives existence, uniqueness and stability by combining a weak notion of contraction with order-theoretic and topological constructs. Doing so frees us from the more restrictive compactness and uniform mixing assumptions found in Hopenhayn and Prescott (1992).

Some of the benefits of weakening these assumptions were discussed above. To put these ideas in a more applied light, suppose that we have a model with unbounded shocks, and, as a result, the state space is unbounded. It is possible to truncate these shocks, thereby creating a version of the model with a compact state space. One immediate problem is that we are approximating in an ad hoc manner, and this approximation may change qualitative and quantitative features of the model. A second problem is that the stability problem might now be significantly harder, because we have reduced the amount of mixing in the model.² A third problem is that estimation might be more difficult because the shock distribution, which determines the likelihood function, has been transformed from a standard to a nonstandard distribution. A fourth problem is that certain questions become more difficult to address, such as whether large shocks are destabilizing, or whether the tails of the stationary distribution have certain properties. For all of these reasons it may be preferable to work with the original model. As we show below, this can be done in a natural and convenient way.

Our results are illustrated in two applications: a model of renewable resource exploitation and an overlapping generations model of the wealth distribution. In both applications, we illustrate situations where the conditions of our theorem are satisfied while those of previous results are not. In fact no current theory from the

²As an example of how truncation might interfere with mixing, consider a model where, in the absence of shocks, the dynamics would yield multiple locally stable steady states (see, e.g., Azariadis and Drazen, 1990). When shocks are present, these points will still be locally attracting "on average." Unless the shocks are sufficiently large, the state might not be able to escape from their basin of attraction. (Stokey, Lucas and Prescott (1989, p. 381) provide an intuitive description of this idea.) In this case initial conditions have permanent effects, and global stability fails because of insufficient mixing.

literature on Markov processes can be used to obtain stability in these cases.

Concerning related literature, the stability of monotone economic models with the Markov property has been studied by Razin and Yahav (1979), Stokey, Lucas and Prescott (1989), Hopenhayn and Prescott (1992), Bhattacharya and Majumdar (2001) and Szeidl (2012). Studies of monotone Markov theory in the mathematical literature include Dubins and Freedman (1966), Yahav (1975), Bhattacharya and Lee (1988), Heikkila and Salonen (1996), Chueshov (2002) and Bhattacharya *et al.* (2010).

The studies of Yahav (1975), Razin and Yahav (1979), Stokey, Lucas and Prescott (1989) and Hopenhayn and Prescott (1992) all use a certain monotone mixing condition suitable for compact, order bounded state spaces. As clarified below, our order reversing condition is weaker than this monotone mixing condition. The papers by Dubins and Freedman (1966), Bhattacharya and Lee (1988), Bhattacharya and Majumdar (2001) and Bhattacharya *et al.* (2010) analyze stability in the monotone setting via a mixing condition called “splitting.” Our order reversing condition is also weaker than splitting. At the same time, the literature on splitting contains important results not treated in this paper.

The paper by Szeidl (2012) is, like our paper, a direct extension of the Hopenhayn-Prezcott stability results for monotone economies. It studies processes that satisfy a certain “weak mixing” condition. Our order reversing condition is weaker than this weak mixing condition, and the main stability results in Szeidl’s paper are special cases of theorems 3.1 and 3.2 below. Nonetheless, Szeidl’s paper contains many thoughtful arguments, and his weak mixing condition can be viewed as a useful way to establish our concept of order reversing.

The work by Chueshov (2002) is a contribution to random dynamical systems theory. It permits unbounded state spaces, but requires continuity throughout, and uses a set of sufficient conditions not directed towards economic applications. Finally, Heikkila and Salonen (1996) provide some extensions to the existence component of Hopenhayn and Prescott’s results that are applicable in non-compact state spaces, but do not treat global stability.

The rest of the paper is structured as follows: Section 2 reviews some basic definitions and introduces the concept of order reversing. Section 3 states the main results and compares them to the existing literature. Section 4 gives applications and section 5 concludes. Proofs can be found in section 6.

2 Preliminaries

At each time $t = 0, 1, \dots$, the state of the economy is described by a point X_t in topological space S . The space S is equipped with its Borel sets \mathcal{B}_S and a closed partial order \leq . An order interval of S is a set of the form $[a, b] := \{x \in S : a \leq x \leq b\}$. A function $f: S \rightarrow \mathbb{R}$ is called *increasing* if $f(x) \leq f(y)$ whenever $x \leq y$. A subset B of S is called *order bounded* if there exists an order interval $[a, b] \subset S$ with $B \subset [a, b]$. In addition, B is called *increasing* if its indicator function $\mathbb{1}_B$ is increasing, and *decreasing* if $\mathbb{1}_B$ is decreasing.

To simplify terminology, we often use the word “distribution” to mean “probability measure on (S, \mathcal{B}_S) ”. The set of all probability measures on (S, \mathcal{B}_S) will be denote by \mathcal{P}_S . We let cbS denote the continuous bounded functions from S to \mathbb{R} , and ibS denote the set of increasing bounded measurable functions from S to \mathbb{R} . We adopt the standard definitions of convergence in distribution and stochastic domination: Given sequence $\{\mu_n\}_{n=0}^\infty$ in \mathcal{P}_S , we say that μ_n *converges to* μ and write $\mu_n \rightarrow \mu_0$ if $\int h d\mu_n \rightarrow \int h d\mu_0$ for all $h \in cbS$. We say that μ_2 *stochastically dominates* μ_1 and write $\mu_1 \preceq \mu_2$ if $\int h d\mu_1 \leq \int h d\mu_2$ for all $h \in ibS$.

Following Hopenhayn and Prescott (1992), we assume that S is a normally ordered Polish space.³ Hopenhayn and Prescott assume in addition that S is compact, with least element a and greatest element b . (A point a is called a *least element* of S if $a \in S$ and $a \leq x$ for all $x \in S$. A point b is called a *greatest element* of S if $b \in S$ and $x \leq b$ for all $x \in S$.) Since we wish to include more general state spaces such as \mathbb{R}^n , we make the weaker assumption that a subset of S is compact if and only if it is closed and order bounded. This is obviously the case in Hopenhayn and Prescott’s setting, where all subsets of S are order bounded, and any closed subset is compact. It also holds for $S = \mathbb{R}^n$ with its standard partial order, since order boundedness is then equivalent to boundedness. In addition, it holds in common state spaces such as \mathbb{R}_+^n or \mathbb{R}_{++}^n , or in any set of the form $I_1 \times \dots \times I_n \subset \mathbb{R}^n$, where each I_i is an open, closed, half-open or half-closed interval in \mathbb{R} .⁴

³A *Polish space* is a separable and completely metrizable topological space. The space (S, \leq) is *normally ordered* if, given any closed increasing set I and closed decreasing set D with $I \cap D = \emptyset$, there exists an f in $ibS \cap cbS$ such that $f(x) = 0$ for all $x \in D$ and $f(x) = 1$ for all $x \in I$.

⁴A simple example that does not satisfy our assumptions is $S = (0, 1) \cup (2, 3)$. In this case the order interval $[0.5, 2.5]$ is closed and order bounded but not compact.

2.1 Markov Properties

Throughout the paper, we suppose that the model under consideration is time-homogeneous and Markovian. The dynamics of such a model can be summarized by a stochastic kernel Q , where $Q(x, B)$ represents the probability that the state moves from $x \in S$ to $B \in \mathcal{B}_S$ in one unit of time. As usual, we require that $Q(x, \cdot) \in \mathcal{P}_S$ for each $x \in S$, and that $Q(\cdot, B)$ is measurable for each $B \in \mathcal{B}_S$. For each $t \in \mathbb{N}$, let Q^t be the t -th order kernel, defined by

$$Q^1 := Q, \quad Q^t(x, B) := \int Q^{t-1}(y, B)Q(x, dy) \quad (x \in S, B \in \mathcal{B}_S).$$

The value $Q^t(x, B)$ represents the probability of transitioning from x to B in t steps.

Here and below, $(\Omega, \mathcal{F}, \mathbb{P})$ denotes a fixed probability space on which all random variables are defined, and \mathbb{E} is the corresponding expectations operator. Given $\mu \in \mathcal{P}_S$ and stochastic kernel Q , an S -valued stochastic process $\{X_t\}_{t \in \mathbb{Z}_+}$ is called (Q, μ) -Markov if X_0 has distribution μ and $Q(x, \cdot)$ is the conditional distribution of X_{t+1} given $X_t = x$.⁵ If μ is the distribution $\delta_x \in \mathcal{P}_S$ concentrated on $x \in S$, we call $\{X_t\}$ (Q, x) -Markov. We call $\{X_t\}$ Q -Markov if $\{X_t\}$ is (Q, μ) -Markov for some $\mu \in \mathcal{P}_S$.

Example 2.1. Many economic models result in processes for the state variables represented by nonlinear, vector-valued stochastic difference equations. As a generic example, consider the S -valued process

$$X_{t+1} = F(X_t, \xi_{t+1}), \quad \{\xi_t\} \stackrel{\text{i.i.d.}}{\sim} \phi, \quad (1)$$

where $\{\xi_t\}$ takes values in $Z \subset \mathbb{R}^m$, the function $F: S \times Z \rightarrow S$ is measurable, and ϕ is a probability measure on the Borel sets of Z . Let Q_F be the kernel

$$Q_F(x, B) := \mathbb{P}\{F(x, \xi_t) \in B\} = \phi\{z \in Z : F(x, z) \in B\}. \quad (2)$$

Then $\{X_t\}$ in (1) is Q_F -Markov.⁶

For each Q we define two operators, sometimes called the left and right Markov operators. The left Markov operator maps $\mu \in \mathcal{P}_S$ into $\mu Q \in \mathcal{P}_S$, where

$$(\mu Q)(B) := \int Q(x, B)\mu(dx) \quad (B \in \mathcal{B}_S). \quad (3)$$

⁵More formally, $\mathbb{P}[X_{t+1} \in B | \mathcal{F}_t] = Q(X_t, B)$ almost surely for all $B \in \mathcal{B}_S$, where \mathcal{F}_t is the σ -algebra generated by the history X_0, \dots, X_t .

⁶Although the process (1) is only first order, models including higher order lags of the state and shock process can be rewritten in the form of (1) by redefining the state variables.

The right Markov operator maps bounded measurable function $h: S \rightarrow \mathbb{R}$ into bounded measurable function Qh , where

$$(Qh)(x) := \int h(y)Q(x, dy) \quad (x \in S).$$

The interpretation of the left Markov operator $\mu \mapsto \mu Q$ is that it shifts the distribution for the state forward by one time period. In particular, if $\{X_t\}$ is (Q, μ) -Markov, then μQ^t is the distribution of X_t . The interpretation of the right Markov operator $h \mapsto Qh$ is that $(Q^t h)(x)$ is the expectation of $h(X_t)$ given $X_0 = x$. If Q_F is the kernel in (2), then $(Q_F h)(x) = \int h[F(x, z)]\phi(dz)$. Also, given any $x \in S$, $B \in \mathcal{B}_S$ and $t \in \mathbb{N}$, the t -th order kernel and the left and right Markov operators are related by $Q^t(x, B) = (\delta_x Q^t)(B) = (Q^t \mathbb{1}_B)(x)$. Here $\mathbb{1}_B$ is the indicator function of B .

A sequence $\{\mu_n\} \subset \mathcal{P}_S$ is called *tight* if, for all $\epsilon > 0$, there exists a compact $K \subset S$ such that $\mu_n(K \setminus S) \leq \epsilon$ for all n . A stochastic kernel Q is called *bounded in probability* if the sequence $\{Q^t(x, \cdot)\}_{t \geq 0}$ is tight for all $x \in S$. If $\mu^* \in \mathcal{P}_S$ and $\mu^* Q = \mu^*$, then μ^* is called *stationary* (or *invariant*) for Q . If Q has a unique stationary distribution μ^* in \mathcal{P}_S , and, in addition, $\mu Q^t \rightarrow \mu^*$ as $t \rightarrow \infty$ for all $\mu \in \mathcal{P}_S$, then Q is called *globally stable*. In this case, μ^* is naturally interpreted as the long-run equilibrium of the economic system. If μ^* is stationary, then any (Q, μ^*) -Markov process $\{X_t\}$ is strict-sense stationary with $X_t \sim \mu^*$ for all t .

If $\mu \in \mathcal{P}_S$ and $\mu Q \preceq \mu$, then μ is called *excessive*. If $\mu \preceq \mu Q$, then μ is called *deficient*. If Q satisfies $\mu Q \preceq \mu' Q$ whenever $\mu \preceq \mu'$, then Q is called *increasing*.⁷ It is in fact sufficient to check that $Q(x, \cdot) \preceq Q(x', \cdot)$ whenever $x \leq x'$. A third equivalent condition is that $Qh \in ibS$ whenever $h \in ibS$. If, on the other hand, $Qh \in cbS$ whenever $h \in cbS$, then Q is called *Feller*.

Remark 2.1. Let Q be an increasing stochastic kernel. If A is an increasing set, then $x \mapsto Q(x, A)$ is increasing. If A is a decreasing set, then $x \mapsto Q(x, A)$ is decreasing.

Remark 2.2. If S has a least element a , then δ_a is deficient for any kernel Q , because $\delta_a \preceq \mu$ for every $\mu \in \mathcal{P}_S$, and hence $\delta_a \preceq \delta_a Q$. Similarly, if S has a greatest element b , then δ_b is excessive for Q .

⁷Many examples of models with increasing kernels were given in the introduction. Other examples not discussed there include various infinite horizon optimal growth models with features such as irreversible investment, renewable resources, distortions, and capital-dependent utility. Increasing kernels are also found in stochastic OLG models besides those mentioned previously, such as models with limited commitment, and in a variety of stochastic games. See, for example, Olson (1989), Amir (2002), Gong et al. (2010), Balbus et al. (2010), and Mirman et al. (2008).

Remark 2.3. Let F and Q_F be as in example 2.1. If $x \mapsto F(x, z)$ is increasing, then Q_F is increasing. If $x \mapsto F(x, z)$ is continuous, then Q_F is Feller.

2.2 Order Reversing

Next we introduce our order-theoretic mixing condition. Let Q be a stochastic kernel on S . We call Q *order reversing* if, for any given x and x' in S with $x \geq x'$, and any independent Q -Markov processes $\{X_t\}$ and $\{X'_t\}$ starting at x and x' respectively, there exists a $t \in \mathbb{N}$ with $\mathbb{P}\{X_t \leq X'_t\} > 0$. In other words, there exists a point in time at which the initial ordering is reversed with positive probability.

Example 2.2. Suppose we are studying a model of household wealth dynamics. Informally, the model is order reversing if, for two households receiving idiosyncratic shocks from the same distribution, it is the case that, regardless of the initial ranking of the two households according to wealth, the probability that their relative wealth positions will be reversed at some point in time is strictly positive.

We make three preliminary comments on the definition. First, in verifying order reversing, it is clearly sufficient to check the existence of a t with $\mathbb{P}\{X_t \leq X'_t\} > 0$ for *arbitrary* pair $x, x' \in S$. Often this is just as easy, and much of the following discussion proceeds accordingly. Second, once x and x' are chosen, there are many pairs of independent Q -Markov processes $\{X_t\}$ and $\{X'_t\}$ starting at x and x' respectively, just as there are many random variables having a given distribution F . It is enough to check that there exists a $t \in \mathbb{N}$ with $\mathbb{P}\{X_t \leq X'_t\} > 0$ for any one of these pairs $\{X_t\}$ and $\{X'_t\}$, because all such pairs have the same joint distribution. Third, it is not entirely clear from the definition given above that order reversing is a property of Q alone. This fact is clarified in the technical appendix, where we give an alternative, more formal, definition.

In remark 2.4 below, we show that for any increasing kernel Q , order reversing is weaker than the monotone mixing condition (MMC) used in Hopenhayn and Prescott (1992). For increasing kernels, order reversing is also weaker than the splitting condition used by Bhattacharya and Majumdar (2001), the “weak mixing” condition used by Szeidl (2012), and the “order mixing” condition used by Kamihigashi and Stachurski (2011a). The proofs are quite straightforward, and details are available from the authors.

Remark 2.4. Let S be a compact metric space with least element a and greatest element b , and let Q be an increasing kernel on S . In this setting, Q is said to satisfy the

MMC whenever

$$\exists \bar{x} \in S \text{ and } k \in \mathbb{N} \text{ such that } Q^k(a, [\bar{x}, b]) > 0 \text{ and } Q^k(b, [a, \bar{x}]) > 0. \quad (4)$$

Under these conditions, Q is order reversing: If we start independent Q -Markov processes $\{X_t^a\}$ and $\{X_t^b\}$ at a and b respectively, then (4) implies the order reversal $X_k^b \leq X_k^a$ occurs at time k with positive probability. Since Q is increasing, closer initial conditions only make this event more likely.⁸

Remark 2.5. To see that order reversing is strictly weaker than the MMC, consider the stochastic kernel $Q(x, B) = \mathbb{P}\{\rho x + \zeta_t \in B\}$ on $S = \mathbb{R}$ associated with the linear Gaussian model

$$X_{t+1} = \rho X_t + \zeta_{t+1}, \quad \{\zeta_t\} \stackrel{\text{iid}}{\sim} N(0, 1). \quad (5)$$

The MMC cannot be applied here, because $S = \mathbb{R}$ and hence the state possesses neither a least nor a greatest element. On the other hand, Q is order reversing. To see this, fix $(x, x') \in \mathbb{R}^2$, and take a second Q -Markov process $X'_{t+1} = \rho X'_t + \zeta'_{t+1}$, where $X'_0 = x'$, $X_0 = x$, and $\{\zeta_t\}$ and $\{\zeta'_t\}$ are IID, standard normal, and independent of each other. The condition $\mathbb{P}\{X_t \leq X'_t\} > 0$ is satisfied with $t = 1$, because

$$\mathbb{P}\{X_1 \leq X'_1\} = \mathbb{P}\{\rho x + \zeta_1 \leq \rho x' + \zeta'_1\} = \mathbb{P}\{\zeta_1 - \zeta'_1 \leq \rho(x' - x)\}.$$

Since $\zeta_1 - \zeta'_1$ is Gaussian, this probability is strictly positive.

3 Results

We can now state our main results, which concern stability of increasing, order reversing stochastic kernels.

3.1 Global Stability

Our first result extends Hopenhayn and Prescott's stability theorem to a broader class of models. It also characterizes the set of increasing order reversing kernels that are globally stable. The proof is in section 6.

⁸To be precise, let \bar{x} and k be as in (4). Fix $x, x' \in S$ and let $\{X_t\}$ and $\{X'_t\}$ be independent, (Q, x) -Markov and (Q, x') -Markov respectively. By independence and $\{X_k \leq \bar{x} \leq X'_k\} \subset \{X_k \leq X'_k\}$, we have $\mathbb{P}\{X_k \leq \bar{x}\}\mathbb{P}\{\bar{x} \leq X'_k\} = \mathbb{P}\{X_k \leq \bar{x} \leq X'_k\} \leq \mathbb{P}\{X_k \leq X'_k\}$. But $\mathbb{P}\{\bar{x} \leq X'_k\} = Q^k(x, [a, \bar{x}])$ and $\mathbb{P}\{X_k \leq \bar{x}\} = Q^k(x, [\bar{x}, b])$ are strictly positive by (4) and remark 2.1. Hence Q is order reversing.

Theorem 3.1. *Let Q be a stochastic kernel on S that is both increasing and order reversing. Then Q is globally stable if and only if*

1. Q is bounded in probability, and
2. Q has either a deficient or an excessive distribution.

Remark 3.1. In terms of sufficient conditions for global stability, the order reversing assumption cannot be omitted, even for *existence* of a stationary distribution. In particular, there exist increasing kernels that are bounded in probability and possess an excessive or deficient distribution, but have no stationary distribution.⁹ On the other hand, regarding necessity, neither monotonicity nor order reversal are used in the proof. Global stability alone implies conditions 1–2.

To see that the conditions of theorem 3.1 are weaker than those of Hopenhayn and Prescott’s stability theorem (Hopenhayn and Prescott, 1992, theorem 2), suppose as they do that S is a compact metric space with least element a and greatest element b , and Q is an increasing kernel satisfying the MMC. The conditions of theorem 3.1 then hold. First, Q is increasing by assumption. Second, Q is order reversing, as shown in remark 2.4. Third, Q is bounded in probability, since S is compact and hence $\{Q^t(x, \cdot)\}$ is always tight. Fourth, Q has a deficient distribution because S has a least element (see remark 2.2).

To see that the conditions of theorem 3.1 are strictly weaker than those of Hopenhayn and Prescott, consider the linear Gaussian model (5) with $\rho \in [0, 1)$. Here the Gaussian shocks force us to choose the state space $S = \mathbb{R}$, which is not compact, and the Hopenhayn-Prescott theorem in its original formulation cannot be applied. On the other hand, all the conditions of theorem 3.1 are satisfied.¹⁰ (Of course this is an extremely simple example. Nontrivial applications are presented in section 4.)

Regarding the proof of theorem 3.1, boundedness in probability and existence of an excessive or deficient distribution generalize Hopenhayn and Prescott’s assumption that S is compact and has a least and greatest element. As Hopenhayn and

⁹An example is the kernel Q associated with the deterministic process on $S = \mathbb{R}_+$ defined by $X_{t+1} = 1/2 + \sum_{n=0}^{\infty} \mathbb{1}\{n \leq X_t < n+1\}(n + (X_t - n)/2)$. It is easy to check that $X_{t+1} > X_t$ with probability one, and hence X_{t+1} and X_t can never have the same distribution. On the other hand, Q is increasing, bounded in probability (because each interval $[n, n+1)$ is absorbing) and has the deficient distribution δ_0 (cf., remark 2.2).

¹⁰That the model is order reversing was shown in remark 2.4. Monotonicity follows from remark 2.3. Boundedness in probability is shown below. For existence of a μ with $\mu \preceq \mu Q$, we can take $\mu = N(0, (1 - \rho^2)^{-1})$.

Prescott show, if S is compact and has a least and greatest element, then the Knaster-Tarski fixed point theorem implies that every increasing stochastic kernel has a stationary distribution. Adding the MMC then yields uniqueness and global stability. In our setting, the same arguments cannot be applied. As remark 3.1 shows, our mixing condition is needed even for existence. Our proof of theorem 3.1 is more akin to a contraction mapping argument than to the Knaster-Tarski fixed point theorem.

We make two final comments. First, even when the conditions of theorem 3.1 hold, they may not be trivial to verify. In section 3.2 we provide a variety of techniques for checking the conditions. Further illustration is given in the applications. Second, there is no continuity requirement in theorem 3.1. However, in many applications the kernel Q will have the Feller property (see remark 2.3). If Q is Feller, then condition 2 can be omitted. Since this result is likely to be useful, we state it as a second theorem.

Theorem 3.2. *Let Q be increasing, order reversing, and Feller. Then Q is globally stable if and only if Q is bounded in probability.*

3.2 Verifying the Conditions

Theorem 3.1 requires that Q is increasing, order reversing, bounded in probability, and possesses an excessive or deficient distribution. A sufficient condition for Q to be increasing was given in remark 2.3. In this section, we present a number of sufficient conditions for the remaining properties.

3.2.1 Checking Boundedness in Probability

Boundedness in probability is a standard condition in the Markov process literature. As is well known, if Q is a stochastic kernel on either $S = \mathbb{R}^n$ or $S = \mathbb{R}_+^n$, then Q is bounded in probability whenever $\sup_t \mathbb{E} \|X_t\| < \infty$ for any (Q, x) -Markov process $\{X_t\}$. (The norm $\|\cdot\|$ can be any norm on \mathbb{R}^n .) For example, it is easy to show by this method that the process (5) is bounded in probability whenever $|\rho| < 1$. More systematic approaches to establishing boundedness in probability can be found in Meyn and Tweedie (2009, chapter 12).

3.2.2 Finding Excessive and Deficient Distributions

Condition 2 of theorem 3.1 requires existence of either an excessive or a deficient distribution. If S has a least element or a greatest element then the condition always holds (see remark 2.2). However, there are many settings where S has neither ($S = \mathbb{R}^n$ and $S = \mathbb{R}_{++}^n$ are obvious examples), and the existence is harder to verify. In this case, one can work more carefully with the definition of the model to construct excessive and deficient distributions. One example is Zhang (1997), who constructs such distributions for the stochastic optimal growth model. However, it is useful to have a more systematic method that is relatively straightforward to check in different applications. To this end we provide the following result. In the result, the statement $Q \preceq Q'$ means that $\mu Q \preceq \mu Q'$ for all $\mu \in \mathcal{P}_S$.

Proposition 3.1. *Let Q be a stochastic kernel on S . If there exists another kernel Q' such that Q' is Feller, bounded in probability and $Q \preceq Q'$ (resp., $Q' \preceq Q$), then Q has an excessive (resp., deficient) distribution.*

An illustration of how the proposition can be used is given in section 4.1.

3.2.3 Checking the Order Reversing Property

In this section we give sufficient conditions for order reversing. To state them, we introduce two new definitions: We call kernel Q on S *upward reaching* if, given any (Q, x) -Markov process $\{X_t\}$ and c in S , there exists a $t \in \mathbb{N}$ such that $\mathbb{P}\{X_t \geq c\} > 0$. We call Q *downward reaching* if, given any (Q, x) -Markov process $\{X_t\}$ and c in S , there exists a $t \in \mathbb{N}$ such that $\mathbb{P}\{X_t \leq c\} > 0$. For example, the linear Gaussian process in (5) is both upward and downward reaching: If we fix x, c in $S = \mathbb{R}$ and take $t = 1$, then $\mathbb{P}\{X_1 \leq c\} = \mathbb{P}\{\rho x + \xi_1 \leq c\} = \mathbb{P}\{\xi_1 \leq c - \rho x\}$. This term is strictly positive because the support of ξ_t is all of \mathbb{R} . Hence Q is downward reaching. The proof of upward reaching is similar.

Proposition 3.2. *Suppose that Q is bounded in probability. If Q is either upward or downward reaching, then Q is order reversing.*

It follows that the statements in theorem 3.1 and theorem 3.2 remain valid if order reversing is replaced by either upward or downward reaching.

4 Applications

We now turn to more substantial applications of the results described above.

4.1 Optimal Exploitation of a Renewable Resource

Consider an elementary model of renewable resource exploitation, where a single planner maximizes $\mathbb{E} \sum_{t=0}^{\infty} \beta^t u(c_t)$ subject to $y_{t+1} = \zeta_t f(y_t - c_t)$. Here y_t is the stock of the resource, c_t is consumption, all variables are nonnegative and $\{\zeta_t\} \stackrel{\text{iid}}{\sim} \phi$. For simplicity, we assume that u is bounded with $u' > 0$, $u'' < 0$, and $u'(0) = \infty$. The growth function f for the resource is assumed to satisfy $f(0) = 0$, $f' > 0$, $f'(0) = \infty$ and $f'(\infty) = 0$. Since f is biologically determined, we do not assume it is concave. To study dynamics, we take y_t as the state variable, and consider the optimal process $y_{t+1} = \zeta_t f(y_t - \sigma(y_t))$, where $\sigma(\cdot)$ is an optimal consumption policy. Let Q be the corresponding stochastic kernel. For the state space we take $S = (0, \infty)$. Zero is deliberately excluded from S so that any stationary distribution on S is automatically non-trivial. Models similar to the one described above have been studied by various authors, including Nishimura and Stachurski (2005), Kamihigashi (2007) and Mitra and Roy (2006).

Regarding the shock process $\{\zeta_t\}$, we permit the occurrence of arbitrarily bad shocks. In the natural resource setting, large negative shocks can take the form of a sudden introduction of pollutants (e.g., oil spills), the arrival of invasive species, disease, extreme droughts, earthquakes, fires, storms and floods. Such low-probability events can have catastrophic environmental and financial consequences. The importance of modeling these left-tail events has been highlighted in a number of recent studies, including Clarke and Reed (1994), Yin and Newman (1996), Brock and Carpenter (2010) and Weitzman (2011). For our purposes, we will assume that $\mathbb{P}\{\zeta_t \leq z\} > 0$ for all $z \in S$, and that $\mathbb{E} \zeta_t < \infty$ and $\mathbb{E}(1/\zeta_t) < \infty$.

For this model, one difficulty for stability analysis is that f is not concave, and hence the optimal policy may be discontinuous. As a result, the stochastic kernel Q is not Feller. Moreover, without additional assumptions, the MMC does not apply, Q is not irreducible, the splitting condition fails, the model is not an expected contraction, the state space is unbounded and the standard Harris recurrence conditions are not satisfied.¹¹ On the other hand, theorem 3.1 can easily be applied. Q is still

¹¹For a discussion of irreducibility and Harris recurrence, see Meyn and Tweedie (2009). On the splitting condition, see, e.g., Bhattacharya and Lee (1988), or Bhattacharya and Majumdar (2001).

increasing and bounded in probability (see, e.g., Nishimura and Stachurski, 2005). Existence of an excessive distribution can be established using proposition 3.1.¹² Moreover, the process is downward reaching (and hence order reversing, cf., proposition 3.2) because if y_0 and \bar{y} in S are given, then

$$\mathbb{P}\{y_1 \leq \bar{y}\} = \mathbb{P}\{\xi_1 f(y_0 - \sigma(y_0)) \leq \bar{y}\} = \mathbb{P}\{\xi_1 \leq \bar{y}/f(y_0 - \sigma(y_0))\} > 0. \quad (6)$$

Hence theorem 3.1 applies, and Q is globally stable.

Figure 1 shows a collection of stationary distributions for $\log y_t$, each one corresponding to a different value of the discount factor β .¹³ For this model, a sudden shift in the optimal harvest policy occurs around $\beta = 0.965$. As a result, a very small difference in the patience of the agent can lead to a large difference in the steady state population of the stock.

4.2 Wealth Distribution Dynamics

Next we consider an OLG model of wealth distribution. Following the existing literature, we introduce persistence in inequality by assuming that old agents provide financial support to their child (cf., e.g., Antunes and Cavalcanti, 2007; Antunes et al., 2008; Cardak, 2004; Couch and Morand, 2005; Lloyd-Ellis, 2000; Lloyd-Ellis and Bernhardt, 2000; Owen and Weil, 1998; Piketty, 1997; and Ranjan, 2001). There are idiosyncratic shocks to endowments and production but no aggregate uncertainty. Unlike much of the literature, we assume that the shocks and hence the state space of the model are unbounded. Permitting unbounded shocks in the wealth distribution allows for the investigation of issues of significant current interest for economists. For example, numerous studies have found that wealth holdings across households are strongly concentrated in the upper tail, and also relatively concentrated in the

¹²Since $f' > 0$ and $f'(\infty) = 0$, we can choose positive constants α, β with $\alpha \mathbb{E} \xi_t < 1$ and $f(x) \leq \alpha x + \beta$. Now take $G(x, z) := z(\alpha x + \beta)$, so that $F(x, z) := z f(x - \sigma(x)) \leq z f(x) \leq G(x, z)$. Letting Q_F and Q_G be the corresponding kernels, the last inequality implies $Q_F \preceq Q_G$. It can be shown that Q_G is both bounded in probability and Feller (for details see the working paper version, Kamihigashi and Stachurski, 2011b), so proposition 3.1 applies.

¹³The utility function is $u(x) = 1 - \exp(-\theta x^\gamma)$ and production is $f(x) = x^\alpha \ell(x)$, where ℓ is the logistic function $\ell(x) = a + (b - a)/(1 + \exp(-c(x - d)))$. The parameters are $a = 1, b = 2, c = 20, d = 1, \theta = 0.5, \gamma = 0.9$ and $\alpha = 0.5$. The discount factor β ranges from 0.945 to 0.99. The shock is lognormal $(-0.1, 0.2)$. For details on the calculations including full justification of consistency, see the working paper version (Kamihigashi and Stachurski, 2011b).




Figure 1: Stationary distributions as a function of β

left tail (see the survey of Davies and Shorrocks, 2000). This leads naturally to modeling with heavy-tailed (and, in particular, unbounded) distributions, such as Pareto or other power law distributions (e.g., Levy and Levy, 2003 or Benhabib *et al.*, 2011). Here we simply assume generic unbounded shocks, and leave the connection to fat tails for future research.

In the model, agents live for two periods and consume only when old. Households consist of one old agent and one child. There is a unit mass of such households indexed by $i \in [0, 1]$. In each period t , the old agent of household i provides financial support b_t^i to her child. The child has the option to become an entrepreneur, investing one unit of the consumption good in a “project,” and receiving stochastic output $\theta + \eta_{t+1}^i$ in period $t + 1$. Let $k_{t+1}^i \in \{0, 1\}$ be young agent i 's investment in the project. If the remainder $b_t^i - k_{t+1}^i$ is positive, then she invests this quantity at the world risk-free rate R . If it is negative then she borrows $k_{t+1}^i - b_t^i$ at the same risk-free rate. Independent of her investment choice, she receives an endowment of e_{t+1}^i units of the consumption good when old. Suppressing the i superscript to simplify notation, her wealth at the beginning of period $t + 1$ is therefore

$$w_{t+1} = (\theta + \eta_{t+1})k_{t+1} - R(k_{t+1} - b_t) + e_{t+1}. \quad (7)$$

We assume that

$$e_{t+1} = \rho e_t + \epsilon_{t+1}, \quad 0 < \rho < 1. \quad (8)$$

The idiosyncratic shocks $\{\eta_t\}$ and $\{\epsilon_t\}$ are taken to be IID and nonnegative, and ϵ_t

satisfies $\mathbb{P}\{\epsilon_t > z\} > 0$ for any $z \geq 0$. (For example, ϵ_t might be lognormal.) We also assume that $R < \theta$, which implies that becoming an entrepreneur is always profitable, even *ex-post*, and every agent would choose to do so absent additional constraint. Due to a credit market imperfection, however, each agent may borrow only up to a fraction $\lambda \in (0, 1)$ of $\theta + \rho e_t$, the minimum possible value of her old-age income (cf., e.g., Matsuyama, 2004). That is,

$$R(k_{t+1} - b_t) \leq \lambda(\theta + \rho e_t). \quad (9)$$

As becoming an entrepreneur is always profitable, young agents do so whenever feasible, implying

$$k_{t+1} = \kappa(b_t, e_t) := \mathbb{1}\{R(1 - b_t) \leq \lambda(\theta + \rho e_t)\}. \quad (10)$$

(Here $\mathbb{1}\{\cdot\}$ is an indicator function.) Let c_{t+1} denote consumption at $t + 1$. It is common in the literature on wealth distribution to assume that each agent derives utility from her own consumption and financial support to her child. Following this approach, we assume that young agents maximize $\mathbb{E}_t[c_{t+1}^{1-\gamma} b_{t+1}^\gamma]$ subject to (7), (9), and the budget constraint $c_{t+1} + b_{t+1} = w_{t+1}$. Regarding the parameter γ we assume that $\gamma R < 1$. Maximization of $c_{t+1}^{1-\gamma} b_{t+1}^\gamma$ subject to the budget constraint implies that $b_{t+1} = \gamma w_{t+1}$. Combining this equality, (7) and (8), we obtain

$$b_{t+1} = \gamma[(\theta + \eta_{t+1} - R)\kappa(b_t, e_t) + Rb_t + \rho e_t + \epsilon_{t+1}]. \quad (11)$$

Together, (8) and (11) define a Markov process with state vector $X_t := (b_t, e_t)$ taking values in state space $S := \mathbb{R}_+^2$. Let Q denote the corresponding stochastic kernel.¹⁴

Recalling that $R < \theta$, $\rho \in (0, 1)$ and $\eta_{t+1} \geq 0$, and observing that $\kappa(b_t, e_t)$ is increasing in (b_t, e_t) , we can see from (8) and (11) that (b_{t+1}, e_{t+1}) is increasing in (b_t, e_t) when the values of the shocks are held fixed. Hence Q is increasing (cf., remark 2.3). On the other hand, (11) is discontinuous in (b_t, e_t) , so Q is not Feller.

As far as we are aware, no existing Markov process theory can be used to show that Q is globally stable unless additional conditions are imposed. In contrast, global stability can be obtained in a straightforward way from theorem 3.1. To begin, let $m_\eta := \mathbb{E} \eta_t$ and $m_\epsilon := \mathbb{E} \epsilon_t$. To see that Q is bounded in probability, we can take expectations of (8) and iterate backwards to obtain

$$\mathbb{E} e_t \leq m_\epsilon / (1 - \rho) + \rho^t e_0 \leq m_\epsilon / (1 - \rho) + e_0 =: \bar{e} \quad (12)$$

¹⁴We do not exclude $(0, 0)$ from the state space since it is not an absorbing state.

for all t . In addition, it follows from (11) and (12) that

$$\mathbb{E} b_{t+1} \leq \gamma[\theta + m_\eta - R + R\mathbb{E} b_t + \bar{e}].$$

Using $\gamma R < 1$ and iterating backwards, we obtain the bound

$$\mathbb{E} b_t \leq \gamma[\theta + m_\eta - R + \bar{e}]/(1 - \gamma R) + b_0 \quad (13)$$

for all t . Together, (12) and (13) imply that Q is bounded in probability.¹⁵ Since $\mathbb{P}\{\epsilon_t > z\} > 0$ for any $z \geq 0$, and since both b_t and e_t can be made arbitrarily large by choosing ϵ_t sufficiently large (see (8) and (11)), it follows that Q is upward reaching, and thus order reversing by proposition 3.2. In view of these results and theorem 3.1, Q will be globally stable whenever it has a deficient or excessive distribution. Since $(0,0)$ is a least element for S , remark 2.2 implies that Q has a deficient distribution, and we conclude that Q is globally stable.

Figure 2 shows smoothed histograms representing the marginal stationary distribution of wealth at two different values of λ , computed by simulation.¹⁶ The shift in the densities shows how the distribution of wealth in the stationary equilibrium can be highly sensitive to the value of the borrowing constraint parameter λ .

5 Conclusion

The methods for analyzing stability of monotone processes developed by Hopenhayn and Prescott (1992) and several other authors have become an important tool in economic modeling. In this paper we introduced a new and very weak mixing condition defined in terms of order, and characterized global stability for monotone models satisfying our condition. Two applications were discussed.

6 Technical Appendix

Before proving theorem 3.1, we need some additional results and notation. To begin, let Q be any stochastic kernel on S , let $x \in S$ and let S -valued stochastic process $\{X_t\}$ be (Q, x) -

¹⁵The function $V(b, e) = |b| + |e|$ is a norm on \mathbb{R}^2 . Equations (12) and (13) yield $\sup_t \mathbb{E}[V(b_t, e_t)] \leq \sup_t \mathbb{E}[b_t] + \sup_t \mathbb{E}[e_t] < \infty$, implying boundedness in probability. See section 3.2.1.

¹⁶The values of λ are 0.57 and 0.58. The other parameters are $\gamma = 0.2$, $R = 1.05$, $\theta = 1.1$ and $\rho = 0.9$. The shock ϵ is lognormal with parameters $\mu = -3$ and $\sigma = 0.1$. The shock η is beta with shape parameters 3,10. For full details on the calculations, see the working paper version (Kamihigashi and Stachurski, 2011b).




Figure 2: Stationary distribution of wealth

Markov. The joint distribution of $\{X_t\}$ over the sequence space S^∞ will be denoted by \mathbf{P}_x^Q . For example, $\mathbf{P}_x^Q\{X_t \in B\} = Q^t(x, B)$ for any $B \subset S$, and $\mathbf{P}_x^Q \cup_{t=0}^\infty \{X_t \in B\}$ is the probability that the process ever enters B . The symbol \mathbf{E}_x^Q represents the expectations operator corresponding to \mathbf{P}_x^Q . For given kernel Q , we say that Borel set $B \subset S$ is

- *strongly accessible* if $\mathbf{P}_x^Q \cup_{t=0}^\infty \{X_t \in B\} = 1$ for all $x \in S$, and
- *C-accessible* if, for all compact $K \subset S$, there exists an $n \in \mathbb{N}$ with $\inf_{x \in K} Q^n(x, B) > 0$.

The following lemma is fundamental to our results, although the proofs are delayed to maintain continuity.

Lemma 6.1. *Let B be a Borel subset of S . If Q is bounded in probability and B is C-accessible, then B is strongly accessible.*

It is helpful to provide a second definition of order reversing. To do so, let

$$\mathbb{G} := \text{graph}(\leq) := \{(y, y') \in S \times S : y \leq y'\},$$

so that $y \leq y'$ iff $(y, y') \in \mathbb{G}$. Also, let Q be a stochastic kernel on S , and consider the product kernel $Q \times Q$ on $S \times S$ defined by

$$(Q \times Q)((x, x'), A \times B) = Q(x, A)Q(x', B) \tag{14}$$

for $(x, x') \in S \times S$ and $A, B \in \mathcal{B}_S$.¹⁷ The product kernel represents the stochastic kernel of the joint process $\{(X_t, X'_t)\}$ when $\{X_t\}$ and $\{X'_t\}$ are independent Q -Markov processes. Using this notation, Q is order reversing if and only if

$$\forall x, x' \in S \text{ with } x' \leq x, \exists t \in \mathbb{N} \text{ such that } (Q \times Q)^t((x, x'), \mathbb{G}) > 0. \quad (15)$$

This second definition emphasizes the fact that order reversing is a property of the kernel Q alone (taking S and \leq as given). Condition (15) can alternatively be written as

$$\forall x, x' \in S \text{ with } x' \leq x, \exists t \geq 0 \text{ such that } \mathbf{P}_{x, x'}^{Q \times Q} \{X_t \leq X'_t\} > 0, \quad (16)$$

where $\{X_t\}$ and $\{X'_t\}$ are independent of each other and (Q, x) -Markov and (Q, x') -Markov respectively. Following Kamihigashi and Stachurski (2011a), Q is called *order mixing* if $\mathbf{P}_{x, x'}^{Q \times Q} \cup_{t=0}^{\infty} \{X_t \leq X'_t\} = 1$ for all $x, x' \in S$. Put differently, Q is order mixing if \mathbb{G} is strongly accessible for the product kernel $Q \times Q$.

Lemma 6.2. *If Q is bounded in probability on S , then so $Q \times Q$ on $S \times S$.*

Lemma 6.3. *If Q is increasing and bounded in probability, then $\{\mu Q^t\}$ is tight for all $\mu \in \mathcal{P}_S$.*

Lemma 6.4. *If Q is increasing and order reversing, then \mathbb{G} is C -accessible for $Q \times Q$.*

Proofs are given at the end of this section.

Let us now turn to the proof of theorem 3.1. The proof proceeds as follows: First we show that under the conditions of the theorem, Q is order mixing. Using order mixing, we then go on to prove existence of a stationary distribution, and global stability.

Lemma 6.5. *If Q is increasing, bounded in probability and order reversing, then Q is order mixing.*

Proof. To show that Q is order mixing we need to prove that \mathbb{G} is strongly accessible for $Q \times Q$ under the conditions of theorem 3.1. Since Q is bounded in probability, $Q \times Q$ is also bounded in probability (lemma 6.2), and hence, by lemma 6.1, it suffices to show that \mathbb{G} is C -accessible for $Q \times Q$. This follows from lemma 6.4. \square

We now prove global stability, making use of order mixing. In the sequel, we define *icbS* to be the bounded, increasing and continuous functions from S to \mathbb{R} (i.e., $icbS = ibS \cap cbS$). To simplify notation, we will also use inner product notation to represent integration, so that

$$\langle \mu, h \rangle := \int h(x) \mu(dx) \quad \text{for } \mu \in \mathcal{P}_S \text{ and } h \in ibS \cup cbS.$$

¹⁷Sets of the form $A \times B$ with $A, B \in \mathcal{B}_S$ provide a semi-ring in the product σ -algebra $\mathcal{B}_S \otimes \mathcal{B}_S$ that also generates $\mathcal{B}_S \otimes \mathcal{B}_S$. Defining the probability measure $Q((x, x'), \cdot)$ on this semi-ring uniquely defines $Q((x, x'), \cdot)$ on all of $\mathcal{B}_S \otimes \mathcal{B}_S$. See, e.g., Dudley (2002, theorem 3.2.7).

It is well known (see, e.g., Stokey *et al.* 1989, p. 219) that the left and right Markov operators are adjoint, in the sense that, for any such h and any $\mu \in \mathcal{P}_S$, we have $\langle \mu, Qh \rangle = \langle \mu Q, h \rangle$.

We will make use of the following results, which are proved at the end of this section.

Lemma 6.6. *Let $\mu, \mu', \mu_n \in \mathcal{P}_S$.*

1. $\mu \preceq \mu'$ iff $\langle \mu, h \rangle \leq \langle \mu', h \rangle$ for all $h \in icbS$,
2. $\mu = \mu'$ iff $\langle \mu, h \rangle = \langle \mu', h \rangle$ for all $h \in icbS$, and
3. $\mu_n \rightarrow \mu$ iff $\{\mu_n\}$ is tight and $\langle \mu_n, h \rangle \rightarrow \langle \mu, h \rangle$ for all $h \in icbS$.

Proof of theorem 3.1. We begin by showing that if Q is globally stable, then conditions 1–2 of the theorem hold. Regarding condition 1, fix $x \in S$. Global stability implies that $\{\mu Q^t\}$ is convergent for each $\mu \in \mathcal{P}_S$, and hence $\{Q^t(x, \cdot)\} = \{\delta_x Q^t\}$ is convergent. Since convergent sequences are tight (Dudley, 2002, proposition 9.3.4) and $x \in S$ was arbitrary, we conclude that Q is bounded in probability, and condition 1 is satisfied. Condition 2 is trivial, because global stability implies existence of a stationary distribution, and every stationary distribution is both deficient and excessive.

Next we show that if Q is increasing, order reversing and conditions 1–2 of theorem 3.1 hold, then Q has at least one stationary distribution. By lemma 6.5, Q is order mixing, and hence, by Kamihigashi and Stachurski (2011a, theorem 3.1), for any ν and ν' in \mathcal{P}_S we have

$$\lim_{t \rightarrow \infty} |\langle \nu Q^t, h \rangle - \langle \nu' Q^t, h \rangle| = 0, \quad \forall h \in ibS. \quad (17)$$

By condition 2 of theorem 3.1 there exists a $\mu \in \mathcal{P}_S$ that is either excessive or deficient. In what follows we will assume it is deficient, since the excessive case only changes the direction of inequalities. Since μ is deficient we have $\mu \preceq \mu Q$. Since Q is increasing, we can iterate on this inequality to establish that the sequence $\{\mu Q^t\}$ is monotone increasing in \preceq . By condition 1 of theorem 3.1 and lemma 6.3, the sequence $\{\mu Q^t\}$ is also tight.

By Prohorov's theorem (Dudley, 2002, theorem 11.5.4), tightness implies existence of a subsequence of $\{\mu Q^t\}$ converging to some $\psi^* \in \mathcal{P}_S$. Since $\{\mu Q^t\}$ is \preceq -increasing, it follows that, for any given $h \in icbS$, the entire sequence $\langle \mu Q^t, h \rangle$ converges up to $\langle \psi^*, h \rangle$. Because $\{\mu Q^t\}$ is tight, part 3 of lemma 6.6 implies that $\mu Q^t \rightarrow \psi^*$.

In addition to $\mu Q^t \rightarrow \psi^*$, we also have $\mu Q^t \preceq \psi^*$ for all $t \geq 0$, because for any $h \in icbS$ and $t \geq 0$ we have

$$\langle \mu Q^t, h \rangle \leq \sup_{t \geq 0} \langle \mu Q^t, h \rangle = \lim_{t \rightarrow \infty} \langle \mu Q^t, h \rangle = \langle \psi^*, h \rangle.$$

The inequality $\mu Q^t \preceq \psi^*$ now follows from part 1 of lemma 6.6.

Next, we claim that $\psi^* \preceq \psi^* Q$. To see this, pick any $h \in icbS$. Since $\mu Q^t \preceq \psi^*$ for all t , and since $Qh \in ibS$,

$$\langle \mu Q^t, Qh \rangle \leq \langle \psi^*, Qh \rangle = \langle \psi^* Q, h \rangle.$$

Using this inequality and the fact that $h \in cbS$, we obtain

$$\langle \psi^*, h \rangle = \lim_{t \rightarrow \infty} \langle \mu Q^{t+1}, h \rangle = \lim_{t \rightarrow \infty} \langle \mu Q^t, Qh \rangle \leq \langle \psi^* Q, h \rangle.$$

Hence $\langle \psi^*, h \rangle \leq \langle \psi^* Q, h \rangle$ for all $h \in icbS$, and $\psi^* \preceq \psi^* Q$ as claimed. Iterating on this inequality we obtain $\psi^* \preceq \psi^* Q^t$ for all t .

To summarize our results so far, we have $\mu Q^t \preceq \psi^* \preceq \psi^* Q \preceq \psi^* Q^t$ for all $t \geq 0$, and hence

$$\langle \mu Q^t, h \rangle \leq \langle \psi^*, h \rangle \leq \langle \psi^* Q, h \rangle \leq \langle \psi^* Q^t, h \rangle \quad \text{for all } h \in icbS.$$

Applying (17), we obtain $\langle \psi^*, h \rangle = \langle \psi^* Q, h \rangle$ for all $h \in icbS$. By lemma 6.6, this implies that $\psi^* = \psi^* Q$. In other words, ψ^* is stationary for Q .

It remains to show that Q is globally stable. Fixing $\nu \in \mathcal{P}_S$ and applying (17) again, we have

$$\langle \nu Q^t, h \rangle \rightarrow \langle \psi^*, h \rangle, \quad \forall h \in ibS. \quad (18)$$

Since $icbS \subset ibS$ and $\{\nu Q^t\}$ is tight (cf., lemma 6.3), this implies that $\nu Q^t \rightarrow \psi^*$ (lemma 6.6, part 3). Finally, uniqueness is also immediate, because if ν is also stationary, then by (18) we have $\langle \nu, h \rangle = \langle \psi^*, h \rangle$ for all $h \in icbS$. By lemma 6.6, we then have $\nu = \psi^*$. \square

Proof of theorem 3.2. Under the conditions of the theorem, Q is order mixing, as proved in lemma 6.5. In addition, boundedness in probability and the Feller property guarantee the existence of a stationary distribution by the Krylov-Bogolubov theorem (Meyn and Tweedie, 2009, proposition 12.1.3 and lemma D.5.3). Given existence of a stationary distribution ψ^* , the proof that Q is globally stable is now identical to the proof of the same claim given for theorem 3.1 (see the preceding paragraph). \square

Proof of proposition 3.1. Suppose that Q' is Feller and bounded in probability with $Q' \preceq Q$. By the Krylov-Bogolubov theorem (Meyn and Tweedie, 2009, proposition 12.1.3 and lemma D.5.3), Q' has at least one stationary distribution μ . For this μ we have $\mu = \mu Q' \preceq \mu Q$. In other words, μ is deficient for Q . A similar argument shows that if Q' is Feller and bounded in probability with $Q \preceq Q'$ then Q has an excessive distribution. \square

Proof of proposition 3.2. Let Q be bounded in probability. Suppose first that Q is upward reaching. Pick any $(x, x') \in S \times S$. Let $\{X_t\}$ and $\{X'_t\}$ be independent, (Q, x) -Markov and (Q, x') -Markov respectively. We need to prove existence of a $k \in \mathbb{N}$ such that $\mathbb{P}\{X_k \leq X'_k\} > 0$. Since Q is bounded in probability, there exists a compact $C \subset S$ with $\mathbb{P}\{X_t \in C\} > 0$ for all $t \geq 0$. Since compact sets are assumed to be order bounded, we can take an order interval $[a, b]$ of S with $C \subset [a, b]$. For this a, b we have $\mathbb{P}\{a \leq X_t \leq b\} > 0$ for all $t \geq 0$. As Q is upward reaching, there is a $k \in \mathbb{N}$ such that $\mathbb{P}\{b \leq X'_k\} > 0$. Using independence, we now have

$$\mathbb{P}\{X_k \leq X'_k\} \geq \mathbb{P}\{X_k \leq b \leq X'_k\} = \mathbb{P}\{X_k \leq b\} \mathbb{P}\{b \leq X'_k\} > 0,$$

as was to be shown. The proof for the downward reaching case is similar. \square

Finally, we complete the proof of all remaining lemmas stated in this section.

Proof of lemma 6.1. Let B be a C -accessible subset of S . To prove the lemma, it suffices to show that $\mathbf{P}_x^Q \cup_t \{X_t \in B\} = 1$ whenever $\{Q^t(x, \cdot)\}$ is tight. To this end, fix $x \in S$, and assume that $\{Q^t(x, \cdot)\}$ is tight. Let $\tau := \inf\{t \geq 0 : X_t \in B\}$. Evidently we have $\cup_{t=0}^{\infty} \{X_t \in B\} = \{\tau < \infty\}$. Thus, we need to show that $\mathbf{P}_x^Q \{\tau < \infty\} = 1$.

Fix $\epsilon > 0$. Since $\{Q^t(x, \cdot)\}$ is tight, there exists a compact set C such that

$$\inf_t \mathbf{P}_x^Q \{X_t \in C\} = \inf_t Q^t(x, C) \geq 1 - \epsilon.$$

Since B is C -accessible, there exists an $n \in \mathbb{N}$ and $\delta > 0$ such that $\inf_{y \in C} Q^n(y, B) \geq \delta$. For $t \in \mathbb{N}$, define $p_t := \mathbf{P}_x^Q \{\tau \leq tn\}$. We wish to obtain a relationship between p_t and p_{t+1} . To this end, note that

$$\begin{aligned} \mathbb{1}\{\tau \leq (t+1)n\} &= \mathbb{1}\{\tau \leq tn\} + \mathbb{1}\{\tau > tn\} \mathbb{1}\{\tau \leq (t+1)n\} \\ &\geq \mathbb{1}\{\tau \leq tn\} + \mathbb{1}\{\tau > tn\} \mathbb{1}\{X_{(t+1)n} \in B\} \\ &\geq \mathbb{1}\{\tau \leq tn\} + \mathbb{1}\{\tau > tn\} \mathbb{1}\{X_{tn} \in C\} \mathbb{1}\{X_{(t+1)n} \in B\}. \end{aligned}$$

Taking expectations yields

$$p_{t+1} \geq p_t + \mathbf{E}_x^Q \mathbb{1}\{\tau > tn\} \mathbb{1}\{X_{tn} \in C\} \mathbb{1}\{X_{(t+1)n} \in B\}.$$

We estimate the last expectation as follows:

$$\begin{aligned} &\mathbf{E}_x^Q \mathbb{1}\{\tau > tn\} \mathbb{1}\{X_{tn} \in C\} \mathbb{1}\{X_{(t+1)n} \in B\} \\ &= \mathbf{E}_x^Q [\mathbb{1}\{\tau > tn\} \mathbb{1}\{X_{tn} \in C\} \mathbf{E}_x^Q [\mathbb{1}\{X_{(t+1)n} \in B\} | \mathcal{F}_{tn}]] \\ &= \mathbf{E}_x^Q [\mathbb{1}\{\tau > tn\} \mathbb{1}\{X_{tn} \in C\} Q^n(X_{tn}, B)] \\ &\geq \mathbf{E}_x^Q \mathbb{1}\{\tau > tn\} \mathbb{1}\{X_{tn} \in C\} \delta \\ &= \mathbf{E}_x^Q (1 - \mathbb{1}\{\tau \leq tn\}) \mathbb{1}\{X_{tn} \in C\} \delta \\ &= \mathbf{E}_x^Q \mathbb{1}\{X_{tn} \in C\} \delta - \mathbf{E}_x^Q \mathbb{1}\{\tau \leq tn\} \mathbb{1}\{X_{tn} \in C\} \delta \\ &\geq (1 - \epsilon) \delta - \mathbf{E}_x^Q \mathbb{1}\{\tau \leq tn\} \delta \\ &= (1 - \epsilon) \delta - p_t \delta. \end{aligned}$$

$$\therefore p_{t+1} \geq p_t + (1 - \epsilon) \delta - p_t \delta = (1 - \delta) p_t + (1 - \epsilon) \delta.$$

The unique, globally stable fixed point of $q_{t+1} = (1 - \delta) q_t + (1 - \epsilon) \delta$ is $1 - \epsilon$, so $1 - \epsilon \leq \lim_{t \rightarrow \infty} p_t = \mathbf{P}_x^Q \{\tau < \infty\} \leq 1$. Since ϵ was arbitrary, we obtain $\mathbf{P}_x^Q \{\tau < \infty\} = 1$. \square

Proof of lemma 6.2. Fix $x, x' \in S$ and $\epsilon > 0$. Since Q is bounded in probability, we can choose compact sets C and C' such that

$$Q^t(x, C) \geq (1 - \epsilon)^{1/2} \quad \text{and} \quad Q^t(x', C') \geq (1 - \epsilon)^{1/2} \quad \text{for all } t.$$

$$\therefore (Q \times Q)^t((x, x'), C \times C') = Q^t(x, C)Q^t(x', C') \geq 1 - \epsilon \quad \text{for all } t.$$

Since $C \times C'$ is compact in the product space, $Q \times Q$ is bounded in probability. \square

Proof of lemma 6.3. Fix $\mu \in \mathcal{P}_S$ and $\epsilon > 0$. Since individual elements of \mathcal{P}_S are tight (Dudley, 2002, theorem 11.5.1), we can choose a compact set $C_\mu \subset S$ with $\mu(C_\mu) \geq 1 - \epsilon$. By assumption, we can take an order interval $[a, b]$ of S with $C_\mu \subset [a, b]$. For this a, b , we have

$$\mu([a, b]^c) = \mu(S \setminus [a, b]) \leq \epsilon. \quad (19)$$

By hypothesis, $\{Q^t(x, \cdot)\}$ is tight for all $x \in S$, so we choose compact subsets C_a and C_b of S with $Q^t(a, C_a) \geq 1 - \epsilon$ and $Q^t(b, C_b) \geq 1 - \epsilon$ for all t . Since $C_a \cup C_b$ is also compact, we can take an order interval $[\alpha, \beta]$ of S with $C_a \cup C_b \subset [\alpha, \beta] \subset S$. We then have $Q^t(a, [\alpha, \beta]) \geq 1 - \epsilon$ and $Q^t(b, [\alpha, \beta]) \geq 1 - \epsilon$ for all t . Letting $I_\alpha := \{x \in S : x \geq \alpha\}$ and $D_\beta := \{x \in S : x \leq \beta\}$, this leads to

$$Q^t(a, I_\alpha) \geq 1 - \epsilon \quad \text{and} \quad Q^t(b, D_\beta) \geq 1 - \epsilon \quad \text{for all } t. \quad (20)$$

In view of remark 2.1 and (20), we have

$$a \leq x \implies Q^t(x, I_\alpha) \geq Q^t(a, I_\alpha) \geq 1 - \epsilon,$$

and, by a similar argument,

$$x \leq b \implies Q^t(x, D_\beta) \geq Q^t(b, D_\beta) \geq 1 - \epsilon.$$

Since $[\alpha, \beta] := \{x \in S : \alpha \leq x \leq \beta\} = I_\alpha \cap D_\beta$, we have

$$Q^t(x, [\alpha, \beta]^c) = Q^t(x, D_\beta^c \cup I_\alpha^c) \leq 2 - Q^t(x, D_\beta) - Q^t(x, I_\alpha).$$

This leads to the estimate

$$a \leq x \leq b \implies Q^t(x, [\alpha, \beta]^c) \leq 2\epsilon. \quad (21)$$

Combining (19) and (21), we now have

$$\begin{aligned} \mu Q^t([\alpha, \beta]^c) &= \int Q^t(x, [\alpha, \beta]^c) \mu(dx) \\ &= \int_{[a, b]} Q^t(x, [\alpha, \beta]^c) \mu(dx) + \int_{[a, b]^c} Q^t(x, [\alpha, \beta]^c) \mu(dx) \\ &\leq \int_{[a, b]} 2\epsilon \mu(dx) + \mu([\alpha, \beta]^c) \leq 3\epsilon. \end{aligned}$$

Since $[\alpha, \beta]$ is compact and t is arbitrary, we conclude that $\{\mu Q^t\}$ is tight. \square

Proof of lemma 6.4. Let C be any compact subset of $S \times S$. We need to prove existence of an $n \in \mathbb{N}$ and $\delta > 0$ such that $(Q \times Q)^n((x, x'), \mathbb{G}) \geq \delta$ whenever $(x, x') \in C$. To do so, we introduce the function

$$\psi_n(x, x') := (Q \times Q)^n((x, x'), \mathbb{G}) = \mathbf{P}_{x, x'}^{Q \times Q} \{X_n \leq X'_n\},$$

where (X_n, X'_n) is $(Q \times Q, (x, x'))$ -Markov. Intuitively, since Q is increasing, the event $\{X_n \leq X'_n\}$ becomes less likely as x rises and x' falls, and hence $\psi_n(x, x')$ is decreasing in x and increasing in x' for each n . A routine argument confirms this is the case.

Since $C \subset S \times S$ is compact, we can take an order interval $[a, b]$ of S with $C \subset [a, b] \times [a, b]$.¹⁸ Moreover, since Q is order reversing, we can take an $n \in \mathbb{N}$ such that $\delta := \psi_n(b, a) > 0$. Observe that

$$(x, x') \in C \implies (x, x') \in [a, b] \times [a, b] \implies x \leq b \text{ and } x' \geq a.$$

$$\therefore (x, x') \in C \implies (Q \times Q)^n((x, x'), \mathbb{G}) = \psi_n(x, x') \geq \psi_n(b, a) = \delta.$$

In other words, \mathbb{G} is C -accessible for $Q \times Q$. □

Proof of lemma 6.6. The statement $\mu \preceq \mu'$ iff $\langle \mu, h \rangle \leq \langle \mu', h \rangle$ for all $h \in icbS$ holds for every normally ordered space, as shown by Whitt (1980, theorem 2.6). Moreover, since \preceq is a partial order on \mathcal{P}_S (Kamae and Krengel, 1978, theorem 2), and hence antisymmetric, it follows that $\mu = \mu'$ iff $\langle \mu, h \rangle = \langle \mu', h \rangle$ for all $h \in icbS$. Regarding the third assertion of the lemma, observe first that if $\mu_n \rightarrow \mu$, then since S is Polish the sequence $\{\mu_n\}$ is tight (Dudley, 2002, theorem 11.5.3). The statement $\langle \mu_n, h \rangle \rightarrow \langle \mu, h \rangle$ whenever $h \in icbS$ is obvious. To prove the converse, suppose that $\{\mu_n\}$ is tight and $\langle \mu_n, h \rangle \rightarrow \langle \mu, h \rangle$ for all $h \in icbS$. Take any subsequence $\{\mu_n\}_{n \in \mathbb{N}_1}$ of $\{\mu_n\}$. By tightness and Prohorov's theorem (Dudley, 2002, theorem 11.5.4), this subsequence has a subsubsequence converging to some $\nu \in \mathcal{P}_S$:

$$\exists \mathbb{N}_2 \subset \mathbb{N}_1 \text{ such that } \lim_{n \in \mathbb{N}_2} \langle \mu_n, h \rangle = \langle \nu, h \rangle \text{ for all } h \in cbS.$$

Since $\langle \mu_n, h \rangle \rightarrow \langle \mu, h \rangle$ for all $h \in icbS$, we now have $\lim_{n \in \mathbb{N}_2} \langle \mu_n, h \rangle = \langle \nu, h \rangle = \langle \mu, h \rangle$ for all $h \in icbS$, and hence $\nu = \mu$. We have now shown that every subsequence of $\{\mu_n\}$ has a subsubsequence converging to μ , and hence the entire sequence also converges to μ . □

¹⁸To see this, let K be a compact subset of S with $C \subset K \times K$. (Such a K can be obtained by projecting C onto the first and second axis, and defining K as the union of these projections.) Since K is order bounded in S by assumption, we just choose $a, b \in S$ with $K \subset [a, b]$.

References

- [1] Aghion, Philippe and Patrick Bolton (1997): "A Theory of Trickle-Down Growth and Development," *Review of Economic Studies*, 64, 151–172.
- [2] Amir, Rabah (2002): "Complementarity and Diagonal Dominance in Discounted Stochastic Games," *Annals of Operations Research* 114, 39–56.
- [3] Antunes, Antonio and Tiago Cavalcanti (2007): "Start up Costs, Limited Enforcement, and the Hidden Economy," *European Economic Review*, 51, 203–224.
- [4] Azariadis, Costas, and Allan Drazen (1990): "Threshold Externalities in Economic Development," *The Quarterly Journal of Economics*, 105 (2), 501–526.
- [5] Balbus, Lukasz, Kevin L. Reffett and Lukasz Woźny (2012): "Stationary Markovian Equilibrium in Altruistic Stochastic OLG Models with Limited Commitment," forthcoming, *Journal of Mathematical Economics*.
- [6] Benhabib, Jess, Alberto Bisin and Shenghao Zhu (2011): "The Distribution of Wealth and Fiscal Policy in Economies with Finitely Lived Agents," *Econometrica*, 79 (1), 123–157.
- [7] Bhattacharya, Rabi N. and Oesook Lee (1988): "Asymptotics of a Class of Markov Processes which are Not in General Irreducible," *Annals of Probability*, 16 (3), 1333–1347.
- [8] Bhattacharya, Rabi N. and Mukul Majumdar (2001): "On a Class of Stable Random Dynamical Systems: Theory and Applications," *Journal of Economic Theory*, 96, 208–229.
- [9] Bhattacharya, Rabi N., Mukul Majumdar, and Nigar Hashimzade (2010): "Limit Theorems for Monotone Markov Processes," *Sankhya*, 72-A (1), 170–190.
- [10] Brock, William A. and Stephen R. Carpenter (2010): "Interacting Regime Shifts in Ecosystems: Implication for Early Warnings," *Ecological Monographs*, 80 (3), 353–367.
- [11] Cabrales, Antonio and Hugo A. Hopenhayn, (1997): "Labor-Market Flexibility and Aggregate Employment Volatility," *Carnegie-Rochester Conference Series on Public Policy* 46, 189–228.
- [12] Hidalgo-Cabrillana, Ana (2009): "Endogenous Capital Market Imperfections, Human Capital, and Intergenerational Mobility," *Journal of Development Economics*, 90, 285–298.
- [13] Cardak, Buly A. (2004): "Ability, Education, and Income Inequality," *Journal of Public Economic Theory*, 6, 239–276.

- [14] Chatterjee, Partha and Malik Shukayev (2010): "A Stochastic Dynamic Model of Trade and Growth: Convergence and Diversification," mimeo, National University of Singapore.
- [15] Chueshov, Igor (2002): *Monotone Random Systems Theory and Applications*, Lecture Notes in Mathematics Vol. 1779, Springer.
- [16] Clarke, Harry R. and William J. Reed (1994): "Consumption/Pollution Tradeoffs in an Environment Vulnerable to Pollution-related Catastrophic Collapse," *Journal of Economic Dynamics and Control*, 18 (5), 991–1010.
- [17] Cooley, Thomas F. and Vincenzo Quadrini (2001): "Financial Markets and Firm Dynamics," *American Economic Review*, 91, 1286–1310.
- [18] Couch, Kenneth A. and Olivier F. Morand (2005): "Inequality, Mobility, and the Transmission of Ability," *Journal of Macroeconomics*, 27, 365–377.
- [19] Das, Satya P. (2006): "Trade, Skill Acquisition and Distribution," *Journal of Development Economics*, 2006, 118–141.
- [20] Davies, James B. and Anthony F. Shorrocks (2000): "The Distribution of Wealth," *Handbook of income distribution*, (A. B. Atkinson and F. Bourguignon) Volume 1, 605–675, Elsevier.
- [21] de Hek, Paul A. (1999): "On Endogenous Growth under Uncertainty," *International Economic Review*, 40, 727–744.
- [22] Díaz, Antonia, Joseph Pijoan-Mas and Jose-Victor Ríos-Rull (2003): "Precautionary Savings and Wealth Distribution under Habit Formation Preferences," *Journal of Monetary Economics*, 50, 1257–1291.
- [23] Dubins, Lester E. and David A. Freedman (1966): "Invariant Probabilities for Certain Markov Processes," *The Annals of Mathematical Statistics*, 37 (4), 837–848.
- [24] Dudley, Richard M. (2002): *Real Analysis and Probability*, Cambridge Studies in Advanced Mathematics No. 74, Cambridge University Press.
- [25] Gong, Liutang, Xiaojun Zhao, Yunhon Yang and Zou Hengfu (2010): "Stochastic Growth with Social-Status Concern: the Existence of a Unique Stationary Distribution," *Journal of Mathematical Economics*, 46, 505–518.
- [26] Heikkila, Seppo and Hannu Salonen (1996): "On the Existence of Extremal Stationary Distributions of Markov Processes," University of Turku working paper No. 66.

- [27] Hopenhayn, Hugo A. (1992): "Entry, Exit, and Firm Dynamics in Long Run Equilibrium," *Econometrica*, 60, 1127–1150.
- [28] Hopenhayn, Hugo A. and Edward C. Prescott (1992): "Stochastic Monotonicity and Stationary Distributions for Dynamic Economies," *Econometrica*, 60, 1387–1406.
- [29] Hopenhayn, Hugo A., Roger Rogerson (1993): "Job Turnover and Policy Evaluation: A General Equilibrium Analysis," *Journal of Political Economy*, 101, 915–938.
- [30] Huggett, Mark (1993): "The Risk-Free Rate in Heterogeneous Agent Economies," *Journal of Economic Dynamics and Control*, 17 953–69.
- [31] Joseph, Gilles and Thomas Weitzblum (2003): "Optimal Unemployment Insurance: Transitional Dynamics vs. Steady State," *Review of Economic Dynamics*, 6, 869–884.
- [32] Kamae, Teturo and Ulrich Krengel (1978): "Stochastic Partial Ordering," *Annals of Probability*, 6, 1044–1069.
- [33] Kamihigashi, Takashi (2007): "Stochastic Optimal Growth with Bounded or Unbounded Utility and Bounded or Unbounded Shocks," *Journal of Mathematical Economics*, 43, 477–500.
- [34] Kamihigashi, Takashi and John Stachurski (2011a): "An Order-Theoretic Mixing Condition for Monotone Markov Chains," *Statistics and Probability Letters*, 82, 262–267.
- [35] Kamihigashi, Takashi and John Stachurski (2011b): "Existence, Stability and Computation of Stationary Distributions: An Extension of the Hopenhayn-Prescott Theorem," RIEB Discussion Paper 2011-32.
- [36] Levy, Moshe and Haim Levy (2003): "Investment Talent and the Pareto Wealth Distribution: Theoretical and Experimental Analysis," *Review of Economics and Statistics*, 85 (3), 709–725.
- [37] Lloyd-Ellis, Huw (2000): "Public Education, Occupational Choice and the Growth-Inequality Relationship," *International Economic Review*, 41, 171–201.
- [38] Lloyd-Ellis, Huw and Dan Bernhardt (2000): "Enterprise, Inequality and Economic Development," *Review of Economic Studies*, 67, 147–168.
- [39] Marcet, Albert, Francesc Obiols-Homs and Philippe Weil (2007): "Incomplete Markets, Labor Supply and Capital Accumulation," *Journal of Monetary Economics*, 54, 2621–2635.
- [40] Matsuyama, Kiminori (2004): "Financial Market Globalization, Symmetry-Breaking, and Endogenous Inequality of Nations," *Econometrica*, 72, 853–884.

- [41] Meyn, Sean and Richard L. Tweedie (2009): *Markov Chains and Stochastic Stability*, 2nd Edition, Cambridge University Press, Cambridge.
- [42] Mirman, Leonard J., Olivier F. Morand and Kevin L. Reffett (2008): "A Qualitative Approach to Markovian Equilibrium in Infinite Horizon Economies with Capital," *Journal of Economic Theory*, 139, 75–98.
- [43] Mitra, Tapan and Santanu Roy (2006): "Optimal Exploitation of Renewable Resources under Uncertainty and the Extinction of Species," *Economic Theory*, 28 (1), 1-23.
- [44] Morand, Olivier F. and Kevin L. Reffett (2007): "Stationary Markovian Equilibrium in Overlapping Generation Models with Stochastic Nonclassical Production and Markov Shocks," *Journal of Mathematical Economics*, 43, 501–522.
- [45] Nishimura, Kazuo and John Stachurski (2005): "Stability of Stochastic Optimal Growth Models: A New Approach," *Journal of Economic Theory*, 122 (1), 100–118.
- [46] Olson, Lars J. (1989): "Stochastic Growth with Irreversible Investment," *Journal of Economic Theory*, 47, 101–129.
- [47] Owen, Ann L. and David N. Weil (1998): "Intergenerational Earnings Mobility, Inequality and Growth," *Journal of Monetary Economics*, 41, 71–104.
- [48] Pijoan-Mas, Joseph (2006): "Precautionary Savings or Working Long Hours?" *Review of Economic Dynamics*, 9, 326–352.
- [49] Piketty, Thomas (1997): "The Dynamics of the Wealth Distribution and the Interest Rate with Credit Rationing," *Review of Economic Studies*, 64, 173–189.
- [50] Ranjan, Prija (2001): "Dynamic Evolution of Income Distribution and Credit-Constrained Human Capital Investment in Open Economies," *Journal of International Economics*, 55, 329–358.
- [51] Razin, Assaf and Joseph A. Yahav (1979): "On Stochastic Models of Economic Growth," *International Economic Review*, 20, 599–604.
- [52] Samaniego, Roberto M. (2006): "Industrial Subsidies and Technology Adoption in General Equilibrium," *Journal of Economic Dynamics and Control*, 30, 1589–1614.
- [53] Samaniego, Roberto M. (2008): "Can Technical Change Exacerbate the Effects of Labor Market Sclerosis?" *Journal of Economic Dynamics and Control*, 32, 497–528.
- [54] Stokey, Nancy L. and Robert E. Lucas, with Edward C. Prescott (1989): *Recursive Methods in Economic Dynamics*, Harvard University Press, Cambridge, MA.

- [55] Szeidl, Adam (2012): "Stable Invariant Distribution in Buffer-Stock Saving and Stochastic Growth Models," mimeo.
- [56] Weitzman, Martin L. (2011): "Fat-tailed Uncertainty in the Economics of Catastrophic Climate Change," *Review of Environmental Economics and Policy*, 5 (2), 275–292.
- [57] Whitt, Ward (1980): "Uniform Conditional Stochastic Order," *Journal of Applied Probability*, 17, 112–123.
- [58] Yahav, Joseph A. (1975): "On a Fixed Point Theorem and its Stochastic Equivalent," *Journal of Applied Probability*, 12 (3), 605–611.
- [59] Yin, Runsheng and David H. Newman (1996): "The Effect of Catastrophic Risk on Forest Investment Decisions," *Journal of Environmental Economics and Management*, 31 (2), 186–197.
- [60] Zhang, Yuzhe (2007): "Stochastic Optimal Growth with a Non-Compact State Space," *Journal of Mathematical Economics*, 43, 115–129.