

comments appreciated

Necessity of Transversality Conditions for Stochastic Problems

Takashi Kamihigashi

RIEB
Kobe University
Rokkodai, Nada, Kobe 657-8501
Japan

Phone/Fax: +81-78-803-7015
E-mail: tkamihig@rieb.kobe-u.ac.jp

November 3, 2000

Abstract

This paper establishes (i) an extension of Michel's (1990, Theorem 1) necessity result to an abstract reduced-form model, (ii) a generalization of the results of Weitzman (1973) and Ekeland and Scheinkman (1986), and (iii) a new result that is useful particularly in the case of homogeneous returns. These results are shown for an extremely general discrete-time reduced-form model that does not assume differentiability, continuity, or concavity, and that imposes virtually no restriction on the state spaces. The three results are further extended to a stochastic reduced-form model. The stochastic extensions are easily accomplished since our deterministic model is so general that the stochastic model is in fact a special case of the deterministic model. We apply our stochastic results to a stochastic reduced-form model with homogeneous returns and a general type of stochastic growth model with CRRA utility.

Keywords: Transversality condition, stochastic optimization, stochastic reduced-form model, homogeneous returns, stochastic growth model.

JEL Classification Numbers: C61, D90, G12

1 Introduction

Since Shell (1969) and Halkin (1974), necessity of transversality conditions (TVCs) has been an uneasy matter to economic theorists who use infinite-horizon optimization problems. While various results on necessity of TVCs for deterministic continuous-time problems are established in Kamihigashi (2000b), the existing literature does not provide widely applicable results on necessity of TVCs for stochastic problems.

The most standard type of TVC, which we call the standard TVC (STVC), is the condition that the value of optimal stocks at infinity must be zero. The literature on necessity of the STVC dates back at least to Peleg (1970) and Weitzman (1973). For a deterministic concave problem, Weitzman shows that the STVC is necessary if the return functions are nonnegative and if the objective function is always finite. Peleg shows the same result for a special case.¹ For a concave optimal control problem, Michel (1990) studies more general TVCs that an optimal path has to satisfy against a feasible path that does not cause an infinite loss. Michel's results, however, do not directly deal with the STVC.²

While the above results in fact provide characterizations of optimal paths for concave problems, Ekeland and Scheinkman (1986) focus on necessity of the STVC for a possibly non-concave problem. They show that the STVC is necessary if the utility functions satisfy a certain condition and if there exists a summable nonnegative sequence that majorizes the utility functions for all feasible paths.

All the aforementioned results, however, are concerned with deterministic problems. When it comes to stochastic problems, the knowledge on this issue is severely limited. Stochastic versions of Weitzman's theorem are shown by Zilcha (1976) and Takekuma (1992), but they require additional restrictive assumptions.³ Stochastic versions of the results of Michel (1990) and Ekeland and Scheinkman (1986) are not available in the literature.

Among the main results of this paper are a stochastic version of the necessity part of Michel (1990, Theorem 1) and a stochastic version of a generalization of the TVC results of

¹See also Peleg and Ryder (1972) for a similar result.

²See Kamihigashi (2000b) for discussions of related results for continuous-time models.

³See Zilcha (1978) for results specific to an undiscounted stationary model.

Weitzman (1973) and Ekeland and Scheinkman (1986).

This paper extends the main results of Kamihigashi (2000b) to discrete-time problems. Both deterministic and stochastic cases are considered. Our model in the deterministic case is an extremely general reduced-form model. The model is general in that it does not assume differentiability, continuity, or concavity, and in that the state spaces are arbitrary vector spaces instead of finite-dimensional spaces. Because of this generality, our model in the stochastic case, which is a natural extension of the deterministic model, is in fact a special case of the deterministic model.⁴

For the deterministic case, we extend Michel's necessity result to our abstract reduced-form model. We also establish a result that simultaneously generalizes the TVC results of Weitzman (1973) and Ekeland and Scheinkman (1986). This generalization is significant especially because it does not require Ekeland and Scheinkman's assumption that there exists a summable nonnegative sequence that majorizes the utility functions for all feasible paths. In addition we obtain a new result that is useful particularly in the case of homogeneous returns. While similar results are shown in Kamihigashi (2000b) for deterministic continuous-time models, the results in this paper are shown in a substantially more general setting.⁵ Furthermore our deterministic results are extended to the stochastic case. The stochastic extensions are easily accomplished since, as mentioned above, the stochastic model is a special case of the deterministic model.

We follow Ekeland and Scheinkman (1986) in using directional derivatives instead of support prices. Since the aforementioned results except for their result use support prices, those results rely heavily on the separation theorem. As is well-known, the separation theorem for infinite-dimensional spaces requires severe restrictions. This is one of the reasons why the results of Zilcha (1976) and Takekuma (1992) are not easily applicable. By contrast our results do not require such restrictions since we use directional derivatives instead of support prices. In fact we use a generalized type of directional derivative that is well-defined for any

⁴The idea that a stochastic model can be viewed as a deterministic model is used by Yano (1989) to study the comparative statics of the stationary state of a stochastic growth model.

⁵While Kamihigashi (2000b) assumes that the return functions are differentiable, that the optimal path is interior, and that the state spaces are time-invariant and finite-dimensional, none of these assumptions is used in this paper.

real-valued function whose domain lies in a vector space. This allows us to concentrate on conditions directly related to TVCs.

Another feature of our approach is that multidimensional problems are transformed into one dimensional problems. Since a problem with any state spaces is reduced to a problem with a one-dimensional state space, our results are established in extremely general settings without introducing additional complexities.

The rest of the paper is organized as follows. Section 2 establishes our deterministic results. Section 3 extends them to the stochastic case. Section 4 applies our stochastic results to a stochastic reduced-form model with homogeneous returns and to a general type of stochastic growth model with CRRA utility. Section 5 concludes the paper. All the proofs are collected in Appendix A. Section A.1 establishes the most general versions of our results in a one-dimensional setting. The main results of the paper are proved by simple applications of the one-dimensional results.

2 The Deterministic Case

Consider the following problem.

$$(2.1) \quad \left\{ \begin{array}{l} \text{“} \max_{\{x_t\}_{t=0}^{\infty}} \sum_{t=0}^{\infty} V_t(x_t, x_{t+1}) \text{”} \\ \text{s.t. } x_0 = \bar{x}_0, \quad \forall t \in \mathbb{Z}_+, (x_t, x_{t+1}) \in X_t. \end{array} \right.$$

This section assumes the following assumptions.

Assumption 2.1. There exists a sequence of real vector spaces $\{D_t\}_{t=0}^{\infty}$ such that $\bar{x}_0 \in D_0$ and $\forall t \in \mathbb{Z}_+, X_t \subset D_t \times D_{t+1}$.

Assumption 2.2. $\forall t \in \mathbb{Z}_+, V_t : X_t \rightarrow [-\infty, \infty)$.

Assumptions 2.1 and 2.2 impose virtually no restriction on the model. Since each D_t is allowed to be an arbitrary vector space, it may even be a space of random variables. Therefore, though we consider (2.1) as a deterministic problem here, it in fact includes stochastic problems as special cases. We can work with this abstract setting since the only operations we need on D_t are addition and scalar multiplication.

We say that a sequence $\{x_t\}_{t=0}^\infty$ is a *feasible path* if $x_0 = \bar{x}_0$ and $\forall t \in \mathbb{Z}_+, (x_t, x_{t+1}) \in X_t$. Since in applications the objective function is often not guaranteed to be finite or well-defined for all feasible paths, we use Brock's (1970) notion of weak maximality as our optimality criterion. We say that a feasible path $\{x_t^*\}$ is *optimal* if for any feasible path $\{x_t\}$,

$$(2.2) \quad \varliminf_{T \uparrow \infty} \sum_{t=0}^T [V_t(x_t, x_{t+1}) - V_t(x_t^*, x_{t+1}^*)] \leq 0.^6$$

Remark 2.1. Suppose $\sum_{t=0}^\infty V_t(x_t, x_{t+1})$ exists in $[-\infty, \infty)$ for all feasible paths $\{x_t\}$.⁷ Suppose $\sup \sum_{t=0}^\infty V_t(x_t, x_{t+1})$ is finite, where the sup is taken over all feasible paths $\{x_t\}$. Then a feasible path $\{x_t^*\}$ is optimal iff for any feasible path $\{x_t\}$,

$$(2.3) \quad \sum_{t=0}^\infty V_t(x_t, x_{t+1}) \leq \sum_{t=0}^\infty V_t(x_t^*, x_{t+1}^*).$$

Therefore, our optimality criterion coincides with the usual maximization criterion whenever the latter makes sense.

Our optimality criterion applies even when the usual maximization criterion fails. In addition, it is weaker than the similar criterion with $\overline{\lim}$ replacing $\underline{\lim}$ in (2.2). Thus, though weak maximality is rarely directly used in applications, our results apply to virtually any discrete-time problem. The rest of this section assumes the following.

Assumption 2.3. There exists an optimal path $\{x_t^*\}$.

Since we are only interested in necessary conditions for optimality, this assumption imposes no restriction on the model.

For $t \in \mathbb{Z}_+$ and $d \in D_{t+1}$ such that $(x_t^*, x_{t+1}^* + \epsilon d) \in X_t$ for $\epsilon > 0$ sufficiently small, define

$$(2.4) \quad V_{t,2}(x_t^*, x_{t+1}^*; d) = \lim_{\epsilon \downarrow 0} \frac{V_t(x_t^*, x_{t+1}^* + \epsilon d) - V_t(x_t^*, x_{t+1}^*)}{\epsilon}.$$

⁶To be precise, this inequality requires that the left-hand side is well-defined. This means that the left-hand side does not involve expressions like " $\infty - \infty$ " and " $-\infty + \infty$ ". An implication of this requirement is that $\forall t \in \mathbb{Z}_+, V_t(x_t^*, x_{t+1}^*)$ is finite; for otherwise the left-hand side of (2.2) is undefined for $\{x_t\} = \{x_t^*\}$.

⁷Throughout this paper, $\forall i \in \mathbb{Z}_+, \sum_{t=i}^\infty \equiv \lim_{T \uparrow \infty} \sum_{t=i}^T$. Such sums are not to be interpreted as Lebesgue integrals.

The right-hand side is always well-defined, though possibly equal to $-\infty$ or ∞ , even if V_t is nondifferentiable or discontinuous. Note that if V_t is partially differentiable (in an appropriate sense) with respect to the second argument at (x_t^*, x_{t+1}^*) and if $V_{t,2}(x_t^*, x_{t+1}^*)$ denotes the partial derivative, then

$$(2.5) \quad V_{t,2}(x_t^*, x_{t+1}^*; d) = V_{t,2}(x_t^*, x_{t+1}^*)d.$$

Remark 2.2. All the results in this section (Theorems 2.1–2.3) hold even if $\overline{\lim}$ replaces $\underline{\lim}$ in (2.4).⁸

Theorem 2.1. *Assume Assumptions 2.1–2.3. Suppose $\forall t \in \mathbb{Z}_+, X_t$ is convex and V_t is concave. Then*

$$(2.6) \quad \underline{\lim}_{t \uparrow \infty} V_{t,2}(x_t^*, x_{t+1}^*; x_{t+1} - x_{t+1}^*) \leq 0$$

for any feasible path $\{x_t\}$ such that

$$(2.7) \quad \underline{\lim}_{T \uparrow \infty} \sum_{t=0}^T [V_t(x_t, x_{t+1}) - V_t(x_t^*, x_{t+1}^*)] > -\infty,$$

$$(2.8) \quad \forall t \in \mathbb{Z}_+, \exists \epsilon > 0, \quad (x_t^*, x_{t+1}^* + \epsilon(x_{t+1} - x_{t+1}^*)) \in X_t.$$

Theorem 2.1 is a discrete-time version of Kamihigashi (2000b, Corollary 3.1); recall footnote 5. The necessity part of Michel (1990, Theorem 1) shows a similar result for a special case of (2.1) that assumes, among other things, finite-dimensional state spaces. He shows that there exists a sequence of support price vectors $\{p_t\}$ such that $\underline{\lim}_{t \uparrow \infty} p_t(x_t^* - x_t) \leq 0$ for any feasible path $\{x_t\}$ satisfying (2.7). In our case, $V_{t,2}(x_t^*, x_{t+1}^*; x_{t+1} - x_{t+1}^*)$ corresponds to $p_t(x_t^* - x_t)$. Condition (2.8) is needed here for $V_{t,2}(x_t^*, x_{t+1}^*; x_{t+1} - x_{t+1}^*)$ to be well-defined. Except for these differences, Theorem 2.1 generalizes Michel's necessity result to our abstract setting.

The proof of Theorem 2.1 can be summarized in one line as follows.

$$(2.9) \quad \underline{\lim}_{t \uparrow \infty} V_{t,2}(x_t^*, x_{t+1}^*; x_{t+1} - x_{t+1}^*) \leq \underline{\lim}_{t \uparrow \infty} \overline{\lim}_{T \uparrow \infty} \sum_{s=t+1}^T [V_s(x_s^*, x_{s+1}^*) - V_s(x_s, x_{s+1})] \leq 0.$$

⁸See footnote 20 for why this remark is true. See (3.9) for why we use $\underline{\lim}$ instead of $\overline{\lim}$ in (2.4).

The first inequality (without $\lim_{t \uparrow \infty}$) follows from a simple perturbation argument. The second inequality is a consequence of (2.7).⁹

The rest of this section assumes the following.

Assumption 2.4. $\forall t \in \mathbb{Z}_+, \exists \underline{\lambda}_t \in [0, 1], \forall \lambda \in [\underline{\lambda}_t, 1], (x_t^*, \lambda x_{t+1}^*) \in X_t$ and $\forall \tau \geq t + 1, (\lambda x_\tau^*, \lambda x_{\tau+1}^*) \in X_\tau$.

Remark 2.3. Assumption 2.4 holds if $\forall t \in \mathbb{Z}_+, X_t$ is convex and $(x_t^*, 0), (0, 0) \in X_t$.

Assumption 2.4 means that the optimal path can be shifted proportionally downward starting from any period. The assumption is common to the well-known results on the STVC in the literature since the STVC basically means that no gain should be achieved by shifting the optimal path proportionally downward.

For $t \in \mathbb{N}$ and $\lambda \in \mathbb{R} \setminus \{1\}$ with $(\lambda x_t^*, \lambda x_{t+1}^*) \in X_t$, define

$$(2.10) \quad w_t(\lambda) = \frac{V_t(x_t^*, x_{t+1}^*) - V_t(\lambda x_t^*, \lambda x_{t+1}^*)}{1 - \lambda},$$

$$(2.11) \quad \hat{w}_t(\lambda) = \sup_{\tilde{\lambda} \in [\lambda, 1]} w_t(\tilde{\lambda}),$$

where $\hat{w}_t(\lambda)$ is defined only for $\lambda \in [\underline{\lambda}_0, 1]$, where $\underline{\lambda}_0$ is given by Assumption 2.4.

Remark 2.4. Let $t \in \mathbb{N}$. If $V_t(\lambda x_t^*, \lambda x_{t+1}^*)$ is concave in $\lambda \in [\underline{\lambda}_0, 1]$, then $\forall \lambda \in [\underline{\lambda}_0, 1], \hat{w}_t(\lambda) = w_t(\lambda)$.

Theorem 2.2. *Assume Assumptions 2.1–2.4. Suppose*

$$(2.12) \quad \exists \{b_t\}_{t=1}^\infty \subset \mathbb{R}, \exists \lambda \in [\underline{\lambda}_0, 1], \forall t \in \mathbb{N}, \quad \hat{w}_t(\lambda) \leq b_t.$$

Then (i) (2.13) \Rightarrow (2.14) and (ii) (2.15) \Rightarrow (2.16), where (2.13)–(2.16) are given by

$$(2.13) \quad \overline{\lim}_{T \uparrow \infty} \sum_{t=1}^T b_t < \infty,$$

$$(2.14) \quad \overline{\lim}_{t \uparrow \infty} V_{t,2}(x_t^*, x_{t+1}^*; -x_{t+1}^*) \leq 0,$$

$$(2.15) \quad \sum_{t=1}^\infty b_t \text{ exists in } [-\infty, \infty),$$

$$(2.16) \quad \overline{\lim}_{t \uparrow \infty} V_{t,2}(x_t^*, x_{t+1}^*; -x_{t+1}^*) \leq 0.$$

⁹These arguments are used only indirectly in the proof in Appendix A, where Theorem 2.1 is derived from a one-dimensional result.

Theorem 2.2 is a discrete-time version of Kamihigashi (2000b, Theorem 3.3); recall footnote 5. Conclusion (ii) generalizes the necessity part of Weitzman (1973) except that TVC (2.16) is expressed in terms of generalized directional derivatives instead of support prices. Note that (2.12) and (2.15) follow from Weitzman's assumptions that the return functions are concave and that the objective function is finite for all feasible paths.

Conclusion (ii) also generalizes Ekeland and Scheinkman (1986, Proposition 5.1). Their result assumes the existence of a nonnegative sequence $\{f_t\}$ with $\sum_{t=0}^{\infty} f_t < \infty$ such that for *all* feasible paths $\{x_t\}$, $\forall t \in \mathbb{Z}_+, V_t(x_t, x_{t+1}) \leq f_t$. Theorem 2.2 does not require this restrictive assumption. Though Ekeland and Scheinkman do not directly assume (2.12) and (2.15), it is shown in Kamihigashi (2000a) that their assumptions imply (2.12) and (2.15). See Kamihigashi (2000a) for further discussions on Ekeland and Scheinkman's result and approach.

The proof of conclusion (i) is similar to that of Theorem 2.1. It can be summarized in one line as follows.

$$(2.17) \quad \varliminf_{t \uparrow \infty} V_{t,2}(x_t^*, x_{t+1}^*; -x_{t+1}^*) \leq \varliminf_{t \uparrow \infty} \overline{\lim}_{T \uparrow \infty} \sum_{s=t+1}^T \hat{w}_s(\lambda) \leq 0.$$

The proof of conclusion (ii) uses a similar argument.¹⁰

Our last result in this section uses the following assumption.

Assumption 2.5. $\exists \bar{\mu} > 1, \forall \mu \in (1, \bar{\mu}]$, (i) $(x_0^*, \mu x_1^*) \in X_0$, (ii) $\forall t \in \mathbb{N}, (\mu x_t^*, \mu x_{t+1}^*) \in X_t$, (iii) $V_0(x_0^*, \mu x_1^*) > -\infty$, and (iv) $\forall t \in \mathbb{N}, V_t(\mu x_t^*, \mu x_{t+1}^*) \geq V_t(x_t^*, x_{t+1}^*)$.

This assumption means that the optimal path can be shifted proportionally upward ((i) and (ii)) and that such a shift (if sufficiently small) entails a finite loss in period 0 and nonnegative gains in subsequent periods ((iii) and (iv)). The assumption is innocuous at least for standard models with constant-returns-to-scale technology.

Theorem 2.3. *Assume Assumptions 2.1–2.5. Suppose*

$$(2.18) \quad \exists \lambda \in [\lambda_0, 1], \exists \mu \in (1, \bar{\mu}], \exists \theta \geq 0, \forall t \in \mathbb{N}, \quad \hat{w}_t(\lambda) \leq \theta w_t(\mu).$$

Then TVC (2.16) holds.

¹⁰In the actual proof of Theorem 2.2 in Appendix A, Theorem 2.2 is derived from a one-dimensional result.

Theorem 2.3 is similar to Kamihigashi (2000b, Theorem 3.4), but the proofs of these results are quite different. Basically Kamihigashi (Theorem 3.4) uses $\lim_{\mu \downarrow 1} w_t(\mu)$ in (2.18) instead of $w_t(\mu)$ and its proof relies heavily on differentiability and the Euler equation. The proof of Theorem 2.3, on the other hand, verifies (2.12) and (2.15) using (2.18) and Assumption 2.5. Theorem 2.3 is useful particularly in the case of homogenous returns. See Section 4 for applications of the stochastic version of Theorem 2.3.

3 The Stochastic Case

This section extends the results in the preceding section to the stochastic case. Let (Ω, \mathcal{F}, P) be a probability space. Let E denote the associated expectation operator; i.e., $Ez = \int z(\omega)dP(\omega)$ for any random variable $z : \Omega \rightarrow \overline{\mathbb{R}}$. When it is important to make explicit the dependence of z on ω , we write $Ez(\omega)$ instead of Ez . Consider the following problem.

$$(3.1) \quad \left\{ \begin{array}{l} \text{“} \max_{\{x_t\}_{t=0}^{\infty}} \sum_{t=0}^{\infty} Ev_t(x_t(\omega), x_{t+1}(\omega), \omega) \text{”} \\ \text{s.t. } x_0 = \bar{x}_0, \quad \forall t \in \mathbb{Z}_+, (x_t, x_{t+1}) \in X_t. \end{array} \right.$$

For any two sets Y and Z , let $F(Y, Z)$ denote the set of all functions from Y to Z . The following assumption means that x_t is a random variable in a real vector space.

Assumption 3.1. There exists a sequence of real vector spaces $\{B_t\}_{t=0}^{\infty}$ such that $x_0 \in F(\Omega, B_0)$ and $\forall t \in \mathbb{Z}_+, X_t \subset F(\Omega, B_t) \times F(\Omega, B_{t+1})$.

The following assumption simply means that the expression $Ev_t(x_t(\omega), x_{t+1}(\omega), \omega)$ makes sense.

Assumption 3.2. $\forall t \in \mathbb{Z}_+, \forall (y, z) \in X_t$, (i) $\forall \omega \in \Omega, v_t(y(\omega), z(\omega), \omega) \in [-\infty, \infty]$, (ii) the mapping $v_t(y(\cdot), z(\cdot), \cdot) : \Omega \rightarrow [-\infty, \infty]$ is measurable, and (iii) $Ev_t(y(\omega), z(\omega), \omega)$ exists in $[-\infty, \infty)$.

We say that a sequence $\{x_t\}$ is a *feasible path* if $x_0 = \bar{x}_0$ and $\forall t \in \mathbb{Z}_+, (x_t, x_{t+1}) \in X_t$.¹¹

¹¹In stochastic optimization problems, feasible paths are usually required to be adapted to a filtration. Though such a requirement could be added here, it is unnecessary to our results.

We say that a feasible path $\{x_t^*\}$ is *optimal* if for any feasible path $\{x_t\}$,

$$(3.2) \quad \lim_{T \uparrow \infty} \sum_{t=0}^T [Ev_t(x_t(\omega), x_{t+1}(\omega), \omega) - Ev_t(x_t^*(\omega), x_{t+1}^*(\omega), \omega)] \leq 0.$$

At this point, the results in Section 2 can be applied to the present model by defining $V_t : X_t \rightarrow [-\infty, \infty)$ and D_t for $t \in \mathbb{Z}_+$ as follows.

$$(3.3) \quad V_t(x_t, x_{t+1}) = Ev_t(x_t(\omega), x_{t+1}(\omega), \omega),$$

$$(3.4) \quad D_t = F(\Omega, B_t).$$

Note that Assumptions 3.1 and 3.2 imply Assumptions 2.1 and 2.2. Hence the model here can be viewed as a deterministic problem. In what follows, we establish stochastic versions of the results in Section 2 with TVCs expressed in terms of v_t instead of V_t .

Like Section 2, this section assumes the existence of an optimal path $\{x_t^*\}$. For simplicity, for $(x_t, x_{t+1}) \in X_t$, $v_t(x_t, x_{t+1})$ denotes the random variable $v_t(x_t(\cdot), x_{t+1}(\cdot), \cdot) : \Omega \rightarrow [-\infty, \infty)$. For $t \in \mathbb{Z}_+$ and $d \in F(\Omega, B_{t+1})$ such that $(x_t^*, x_{t+1}^* + \epsilon d) \in X_t$ for $\epsilon > 0$ sufficiently small, we define the random variable $v_{t,2}(x_t^*, x_{t+1}^*; d)$ as in (2.4); to be precise,

$$(3.5) \quad v_{t,2}(x_t^*, x_{t+1}^*; d) = \underline{\lim}_{\epsilon \downarrow 0} \frac{v_t(x_t^*, x_{t+1}^* + \epsilon d) - v_t(x_t^*, x_{t+1}^*)}{\epsilon},$$

where $\underline{\lim}_{\epsilon \downarrow 0}$ is applied pointwise (i.e., for each $\omega \in \Omega$).

Remark 3.1. All the results in this section (Theorems 3.1–3.3) hold even if $\overline{\lim}$ replaces $\underline{\lim}$ in (3.5).¹²

Theorem 3.1. *Assume Assumptions 2.3, 3.1, and 3.2. Suppose $\forall t \in \mathbb{Z}_+, X_t$ is convex and $\forall \omega \in \Omega, v_t(\cdot, \cdot, \omega)$ is concave. Then*

$$(3.6) \quad \underline{\lim}_{t \uparrow \infty} Ev_{t,2}(x_t^*, x_{t+1}^*; x_{t+1} - x_{t+1}^*) \leq 0$$

for any feasible path $\{x_t\}$ satisfying (2.7) and the following:

$$(3.7) \quad \forall t \in \mathbb{Z}_+, \exists \epsilon > 0, \quad \zeta_t(\epsilon) \equiv (x_t^*, x_{t+1}^* + \epsilon(x_{t+1} - x_{t+1}^*)) \in X_t, \quad Ev_t(\zeta_t(\epsilon)) > -\infty.$$

¹²See footnote 20 for why this remark is true. See (3.9) for why we use $\underline{\lim}$ in (3.5).

The proof of Theorem 3.1 uses the last inequality in (3.7) in conjunction with the monotone convergence theorem to show

$$(3.8) \quad \forall t \in \mathbb{Z}_+, \quad Ev_{t,2}(x_t^*, x_{t+1}^*; x_{t+1} - x_{t+1}^*) = V_{t,2}(x_t^*, x_{t+1}^*; x_{t+1} - x_{t+1}^*).$$

Given this result, Theorem 3.1 follows immediately from Theorem 2.1.

The rest of this section assumes Assumption 2.4. Stochastic versions of Theorems 2.2 and 2.3 can easily be shown under the following assumption.

$$\textbf{Assumption 3.3. } \forall t \in \mathbb{Z}_+, Ev_{t,2}(x_t^*, x_{t+1}^*; -x_{t+1}^*) \leq V_{t,2}(x_t^*, x_{t+1}^*; -x_{t+1}^*).$$

The above inequality can be expressed as

$$(3.9) \quad E \underline{\lim}_{\epsilon \downarrow 0} \frac{v_t(x_t^*, x_{t+1}^* - \epsilon x_{t+1}^*) - v_t(x_t^*, x_{t+1}^*)}{\epsilon} \leq \underline{\lim}_{\epsilon \downarrow 0} E \frac{v_t(x_t^*, x_{t+1}^* - \epsilon x_{t+1}^*) - v_t(x_t^*, x_{t+1}^*)}{\epsilon}.$$

Of course this holds by Fatou's lemma under the hypothesis of the lemma.¹³ This is why we use $\underline{\lim}$ instead of $\overline{\lim}$ in (2.4) and (3.5).

Remark 3.2. Assumption 3.3 holds (with equality) if $\forall t \in \mathbb{Z}_+, v_t(x_t^*, \lambda x_{t+1}^*)$ is concave in $\lambda \in [\underline{\lambda}_t, 1]$ ¹⁴ and if $\exists \lambda \in [\underline{\lambda}_t, 1], Ev_t(x_t^*, \lambda x_{t+1}^*) > -\infty$, where $\underline{\lambda}_t$ is given by Assumption 2.4.

Remark 3.3. Assumption 3.3 holds if $\forall t \in \mathbb{Z}_+, v_t(x_t^*, \lambda x_{t+1}^*)$ is nonincreasing in $\lambda \in [\underline{\lambda}_t, 1]$ (which is the case in most economic models).

Theorem 3.2. Assume Assumptions 2.3, 2.4, 3.1–3.3, and (2.12). Then (i) (2.13) \Rightarrow (3.10) and (ii) (2.15) \Rightarrow (3.11), where (3.10) and (3.11) are given by

$$(3.10) \quad \underline{\lim}_{t \uparrow \infty} Ev_{t,2}(x_t^*, x_{t+1}^*; -x_{t+1}^*) \leq 0,$$

$$(3.11) \quad \overline{\lim}_{t \uparrow \infty} Ev_{t,2}(x_t^*, x_{t+1}^*; -x_{t+1}^*) \leq 0.$$

¹³More specifically, (3.9) holds by Fatou's lemma if there exist $\tilde{\epsilon} > 0$ and a measurable function $\xi_t : \Omega \rightarrow \mathbb{R}$ such that

$$E\xi_t > -\infty, \quad \forall \epsilon \in (0, \tilde{\epsilon}), \quad \frac{v_t(x_t^*, x_{t+1}^* - \epsilon x_{t+1}^*) - v_t(x_t^*, x_{t+1}^*)}{\epsilon} \geq \xi_t.$$

¹⁴To be precise, by “ $v_t(x_t^*, \lambda x_{t+1}^*)$ is concave in λ ,” we mean that with probability one, $v_t(x_t^*, \lambda x_{t+1}^*)$ is a concave function of λ . Likewise any condition involving random variables is understood to hold with probability one.

Theorem 3.2 is a stochastic version of Theorem 2.2. Conclusion (ii) generalizes the TVC results of Zilcha (1976) and Takekuma (1992) except that our result uses generalized directional derivatives instead of support prices. Our last general result is a stochastic version of Theorem 2.3.

Theorem 3.3. *Under Assumptions 2.3–2.5, 3.1–3.3, and (2.18), TVC (3.11) holds.*

4 Applications

This section continues to consider the stochastic model (3.1) to offer applications of Theorems 3.2 and 3.3. Assumptions 2.3, 2.4, 3.1, and 3.2 are assumed throughout.

4.1 Homogeneous Returns

In many economic models, the return functions are assumed to be homogenous (e.g., Lucas 1988; Rebelo 1991). In stochastic versions of such models, the following assumption holds.

Assumption 4.1. $\exists \alpha \in \mathbb{R} \setminus \{0\}, \forall t \in \mathbb{N}$, for any $\lambda > 0$ such that $(\lambda x_t^*, \lambda x_{t+1}^*) \in X_t$, we have $v_t(\lambda x_t^*, \lambda x_{t+1}^*) = \lambda^\alpha v_t(x_t^*, x_{t+1}^*)$.

(Proposition 4.3 below deals with the case $\alpha = 0$.) Under certain growth conditions, Alvarez and Stokey (1998) shows the basic results of dynamic programming and the necessity of the STVC for deterministic stationary problems with homogeneous return functions. Here we show that with homogeneous returns, the necessity of the STVC can easily be verified without growth conditions. We use the following standard assumptions.

Assumption 4.2. $\forall t \in \mathbb{Z}_+, E v_t(\lambda x_t^*, \lambda x_{t+1}^*)$ is nondecreasing in $\lambda \in [\underline{\lambda}_0, 1]$, where $\underline{\lambda}_0$ is given by Assumption 2.4.

Assumption 4.3. $\forall t \in \mathbb{Z}_+, v_t(x_t^*, \lambda x_{t+1}^*)$ is nonincreasing in $\lambda \in [\underline{\lambda}_t, 1]$.

Remark 4.1. Under Assumption 4.3, TVC (3.11) is written as

$$(4.1) \quad \lim_{t \uparrow \infty} E v_{t,2}(x_t^*, x_{t+1}^*; -x_{t+1}^*) = 0,$$

which is a stochastic version of the STVC.

Proposition 4.1. *Assume Assumptions 2.3, 2.4, 3.1, 3.2, and 4.1–4.3. Suppose*

$$(4.2) \quad -\infty < \sum_{t=1}^{\infty} Ev_t(x_t^*, x_{t+1}^*) < \infty.^{15}$$

Then TVC (4.1) holds.

The proof of Proposition 4.1 uses Theorem 3.2. Condition (4.2) is usually assumed or taken for granted in applied studies. Proposition 4.1 shows that the STVC is necessary in such cases. Condition (4.2), however, is unnecessary under Assumption 2.5, which allows us to apply Theorem 3.3.

Proposition 4.2. *Under Assumptions 2.3–2.5, 3.1, 3.2, and 4.1–4.3, TVC (4.1) holds.*

Propositions 4.1 and 4.2 indicate that for models with homogeneous returns, there is essentially no issue about necessity of the STVC.

4.2 A Stochastic Growth Model

Consider the following problem.

$$(4.3) \quad \begin{cases} \max_{\{x_t\}_{t=0}^{\infty}} & \sum_{t=0}^{\infty} E\beta^t u(g_t(x_t(\omega), x_{t+1}(\omega), \omega)) \\ \text{s.t.} & x_0 = \bar{x}_0, \quad \forall t \in \mathbb{Z}_+, (x_t, x_{t+1}) \in X_t. \end{cases}$$

As in Section 3, $g_t(x_t, x_{t+1})$ denotes the random variable $g_t(x_t(\cdot), x_{t+1}(\cdot), \cdot)$. Various stochastic growth models (e.g., Brock and Mirman (1972)) take the form of (4.3). One may call x_t the capital stock (or the vector of capital stocks) at the beginning of period t and $g_t(x_t, x_{t+1})$ consumption in period t .¹⁶ For simplicity, we assume the following.

Assumption 4.4. (i) $\beta \in (0, 1)$, (ii) $u : \mathbb{R}_+ \rightarrow [-\infty, \infty)$, and (iii) $\exists \alpha \in (-\infty, 1]$,

$$(4.4) \quad u(\cdot) = \begin{cases} \frac{(\cdot)^\alpha}{\alpha} & \text{if } \alpha \neq 0, \\ \ln(\cdot) & \text{if } \alpha = 0. \end{cases}$$

¹⁵The infinite sum exists since $\forall t \in \mathbb{N}, \alpha Ev_t(x_t^*, x_{t+1}^*) \geq 0$. See (A.30).

¹⁶One can apply the results in this section to models with endogenous labor supply such as RBC models (e.g., King, Rebelo, and Plosser, 1988). To do so, one may take the optimal labor path as given and consider the maximization problem over consumption and capital paths.

Assumption 4.5. $\forall t \in \mathbb{Z}_+$, (i) X_t is convex, (ii) $(x_t^*, 0), (0, 0) \in X_t$, (iii) $g_t(0, 0) \geq 0$, (iv) $g_t(x_t^*, x_{t+1}^*) > 0$, and (v) $\forall \omega \in \Omega, g_t(\cdot, \cdot, \omega)$ is concave.

Assumption 4.6. $\forall t \in \mathbb{Z}_+, g_t(x_t^*, \lambda x_{t+1}^*)$ is nonincreasing and continuous in $\lambda \in (0, 1]$.

Remark 4.2. Under Assumptions 4.4–4.6, TVC (3.11) is written as

$$(4.5) \quad \lim_{t \uparrow \infty} \beta^t E[u'(g_t(x_t^*, x_{t+1}^*))g_{t,2}(x_t^*, x_{t+1}^*; -x_{t+1}^*)] = 0,$$

where $g_{t,2}(x_t^*, x_{t+1}^*; -x_{t+1}^*)$ is defined as in (3.5).¹⁷

Proposition 4.3. *Assume Assumptions 2.3, 3.1, 3.2, and 4.4–4.6. Suppose $\alpha = 0$. Then TVC (4.5) holds.*

Proposition 4.4. *Assume Assumptions 2.3, 3.1, 3.2, and 4.4–4.6. Suppose $\alpha \neq 0$ and*

$$(4.6) \quad -\infty < \sum_{t=1}^{\infty} \beta^t E u(g_t(x_t^*, x_{t+1}^*)) < \infty.$$

Then TVC (4.5) holds.¹⁸

The proofs of Propositions 4.3 and 4.4 use Theorem 3.2. Proposition 4.3 shows that the STVC is always necessary in the logarithmic case. Even in the non-logarithmic case, Proposition 4.4 shows that the STVC is guaranteed to be necessary unless one is willing to allow lifetime utility to be infinite at the optimum. Such cases are rare in practice since (4.6) is usually assumed or taken for granted in applied studies.

Even without (4.6), however, Theorem 3.3 can be invoked under Assumption 2.5. Indeed, if $\forall t \in \mathbb{N}, g_t$ is homogenous, then Proposition 4.2 (a consequence of Theorem 3.3) directly applies. We also have a useful result that does not assume homogeneity. The result, whose statement is slightly complicated, is stated in Appendix B.

¹⁷This remark can easily be verified by showing $v_{t,2}(x_t^*, x_{t+1}^*; -x_{t+1}^*) = u'(g_t(x_t^*, x_{t+1}^*))g_{t,2}(x_t^*, x_{t+1}^*; -x_{t+1}^*)$ using the mean value theorem.

¹⁸Proposition 4.4 can easily be generalized to more general utility functions. See footnote 22 for details.

5 Concluding Remarks

In this paper we showed (i) an extension of Michel’s (1990, Theorem 1) necessity result to our abstract reduced-form model, (ii) a generalization of the TVC results of Weitzman (1973) and Ekeland and Scheinkman (1986), and (iii) a new result that is useful particularly in the case of homogeneous returns. These results were shown for an extremely general deterministic reduced-form model that does not assume differentiability, continuity, or concavity, and that imposes virtually no restriction on the state spaces. The three results were further extended to a stochastic reduced-form model. The stochastic extensions were easily accomplished since our deterministic model is so general that the stochastic model is in fact a special case of the deterministic model.

As examples of applications, we studied two special cases of the stochastic model. The first was a stochastic reduced-form model with homogeneous returns. For that model, we showed that the STVC is necessary under standard assumptions, even when the objective function is not guaranteed to be finite at the optimum. The second special case was a general type of stochastic growth model with CRRA utility. For that model, we showed that the STVC is necessary if utility is logarithmic or if the objective function is finite at the optimum.¹⁹

As those special cases illustrate, our general stochastic results are highly useful. They are significant as well not only because the existing literature does not provide widely applicable results on necessity of TVCs for stochastic problems, but also because our stochastic results were established at the same level of generality as that of our very general deterministic results. The results of this paper suggest that as far as necessity of TVCs is concerned, there is little difference between deterministic and stochastic cases.

A Proofs

This appendix proves the results stated in the main text. Section A.1 considers a one-dimensional version of the reduced-form model studied in Section 2. Sections A.2–A.11

¹⁹ Appendix B shows a result that does not require the finiteness of the objective function.

prove the results stated in the main text. All the theorems in Section 2 are derived from the results in Section A.1.

A.1 General Results

Consider the following problem.

$$(A.1) \quad \begin{cases} \max_{\{y_t\}_{t=0}^{\infty}} \sum_{t=0}^{\infty} r_t(y_t, y_{t+1}) \\ \text{s.t. } y_0 = \bar{y}_0, \forall t \in \mathbb{Z}_+, (y_t, y_{t+1}) \in Y_t. \end{cases}$$

Assumption A.1. $\bar{y}_0 \in \mathbb{R}$ and $\forall t \in \mathbb{Z}_+, Y_t \subset \mathbb{R} \times \mathbb{R}$.

Assumption A.2. $r_t : Y_t \rightarrow [-\infty, \infty)$.

Assumption A.1 says that y_t is one-dimensional. Feasible paths and optimal paths are defined as in Section 2.

Assumption A.3. There exists an optimal path $\{y_t^*\}$.

Assumption A.4. $\forall t \in \mathbb{Z}_+, \exists \bar{\epsilon}_t > 0, \forall \epsilon \in (0, \bar{\epsilon}_t], (y_t^*, y_{t+1}^* - \epsilon) \in Y_t$ and $\forall \tau \geq t + 1, (y_\tau^* - \epsilon, y_{\tau+1}^* - \epsilon) \in Y_\tau$.

Assumption A.4 means that the optimal path can be shifted uniformly downward starting from any period. For $t \in \mathbb{Z}_+$ and $d \in \mathbb{R}$, define $r_{t,2}(y_t^*, y_{t+1}^*; d)$ as in (2.4).

Remark A.1. Theorems A.1 and A.2 below hold even if $\overline{\lim}$ replaces $\underline{\lim}$ in (2.4).²⁰

For $t \in \mathbb{Z}_+$, define

$$(A.2) \quad q_t = r_{t,2}(y_t^*, y_{t+1}^*; -1).$$

For $t \in \mathbb{N}$ and $\epsilon \in \mathbb{R} \setminus \{0\}$ with $(y_t^* - \epsilon, y_{t+1}^* - \epsilon) \in Y_t$, define

$$(A.3) \quad m_t(\epsilon) = \frac{r_t(y_t^*, y_{t+1}^*) - r_t(y_t^* - \epsilon, y_{t+1}^* - \epsilon)}{\epsilon},$$

$$(A.4) \quad \hat{m}_t(\epsilon) = \sup_{\tilde{\epsilon} \in (0, \epsilon]} m_t(\tilde{\epsilon}),$$

where $\hat{m}_t(\epsilon)$ is defined only for $\epsilon \in (0, \bar{\epsilon}_0]$, where $\bar{\epsilon}_0$ is given by Assumption A.4.

²⁰This can be seen by replacing $\underline{\lim}$ with $\overline{\lim}$ in the last sentence of the proof of Lemma A.1 and (A.5). Lemma A.1 is the only place where (2.4) is used.

Lemma A.1. Under Assumptions A.1–A.4,

$$(A.5) \quad \forall s \in \mathbb{Z}_+, \quad q_s \leq \underline{\lim}_{\epsilon \downarrow 0} \overline{\lim}_{T \uparrow \infty} \sum_{t=s+1}^T m_t(\epsilon).$$

Proof. Let $s \in \mathbb{Z}_+$ and $\epsilon \in (0, \bar{\epsilon}_s]$. By the optimality of $\{y_t^*\}$ and Assumption A.4,

$$(A.6) \quad \underline{\lim}_{T \uparrow \infty} \left[r_s(y_s^*, y_{s+1}^* - \epsilon) - r_s(y_s^*, y_{s+1}^*) + \sum_{t=s+1}^T [r_t(y_t^* - \epsilon, y_{t+1}^* - \epsilon) - r_t(y_t^*, y_{t+1}^*)] \right] \leq 0.$$

Dividing through by ϵ and rearranging, we get

$$(A.7) \quad \frac{r_s(y_s^*, y_{s+1}^* - \epsilon) - r_s(y_s^*, y_{s+1}^*)}{\epsilon} \leq \overline{\lim}_{T \uparrow \infty} \sum_{t=s+1}^T \frac{r_t(y_t^* - \epsilon, y_{t+1}^* - \epsilon)}{\epsilon}.$$

Recalling (A.3) and applying $\underline{\lim}_{\epsilon \downarrow 0}$ yields (A.5). \square

Define

$$(A.8) \quad \Psi = \{\{f_t\}_{t=1}^\infty \subset \mathbb{R} \mid \overline{\lim}_{T \uparrow \infty} \sum_{t=1}^T f_t \in [-\infty, \infty)\},$$

$$(A.9) \quad \Phi = \{\{f_t\}_{t=1}^\infty \subset \mathbb{R} \mid \lim_{T \uparrow \infty} \sum_{t=1}^T f_t \text{ exists in } [-\infty, \infty)\}.$$

Lemma A.2. (i) $\forall \{f_t\} \in \Psi, \underline{\lim}_{s \uparrow \infty} \overline{\lim}_{T \uparrow \infty} \sum_{t=s}^T f_t \leq 0$.²¹

(ii) $\forall \{f_t\} \in \Phi, \overline{\lim}_{s \uparrow \infty} \overline{\lim}_{T \uparrow \infty} \sum_{t=s}^T f_t \leq 0$.

Proof. Let $\{f_t\} \in \Psi$ and $A = \overline{\lim}_{T \uparrow \infty} \sum_{t=1}^T f_t (< \infty)$. Then

$$(A.10) \quad \forall s \geq 2, \quad \overline{\lim}_{T \uparrow \infty} \sum_{t=s}^T f_t = A - \sum_{t=1}^{s-1} f_t.$$

Thus if $A = -\infty$, then $\underline{\lim}_{s \uparrow \infty} \overline{\lim}_{T \uparrow \infty} \sum_{t=s}^T f_t = -\infty$. If $A > -\infty$, then

$$(A.11) \quad \underline{\lim}_{s \uparrow \infty} \overline{\lim}_{T \uparrow \infty} \sum_{t=s}^T f_t = A - \overline{\lim}_{s \uparrow \infty} \sum_{t=1}^{s-1} f_t = A - A = 0.$$

Hence (i) holds. The proof of (ii) is similar. \square

Theorem A.1. Assume Assumptions A.1–A.4. Suppose

$$(A.12) \quad \exists \{b_t\}_{t=1}^\infty \subset \mathbb{R}, \exists \epsilon \in (0, \bar{\epsilon}_0], \forall t \in \mathbb{N}, \quad \hat{m}_t(\epsilon) \leq b_t.$$

Then (i) $\{b_t\} \in \Psi \Rightarrow \underline{\lim}_{t \uparrow \infty} q_t \leq 0$ and (ii) $\{b_t\} \in \Phi \Rightarrow \overline{\lim}_{t \uparrow \infty} q_t \leq 0$.

²¹A similar result is shown by Michel (1990, Proposition 1). Continuous-time versions of Lemma A.2 and some of the other results in this paper are shown in Kamihigashi (2000b).

Proof. By (A.5) and (A.12), $\forall s \in \mathbb{Z}_+$, $q_s \leq \overline{\lim}_{T \uparrow \infty} \sum_{t=s+1}^T b_t$. Thus both (i) and (ii) hold by Lemma A.2. \square

Assumption A.5. $\exists \bar{\delta} > 0, \forall \delta \in (0, \bar{\delta}]$, (i) $(y_0^*, y_1^* + \delta) \in Y_0$, (ii) $\forall t \in \mathbb{N}, (y_t^* + \delta, y_{t+1}^* + \delta) \in Y_t$, (iii) $r_0(y_0^*, y_1^* + \delta) > -\infty$, and (iv) $\forall t \in \mathbb{N}, r_t(y_t^* + \delta, y_{t+1}^* + \delta) \geq r_t(y_t^*, y_{t+1}^*)$.

Theorem A.2. *Assume Assumptions A.1–A.5. Suppose*

$$(A.13) \quad \exists \epsilon \in (0, \bar{\epsilon}_0], \exists \delta \in (0, \bar{\delta}], \exists \theta \geq 0, \forall t \in \mathbb{N}, \hat{m}_t(\epsilon) \leq \theta m_t(-\delta).$$

Then $\overline{\lim}_{t \uparrow \infty} q_t \leq 0$.

Proof. By the optimality of $\{y_t^*\}$ and Assumption A.5,

$$(A.14) \quad r_0(y_0^*, y_1^* + \delta) - r_0(y_0^*, y_1^*) + \sum_{t=1}^{\infty} [r_t(y_t^* + \delta, y_{t+1}^* + \delta) - r_t(y_t^*, y_{t+1}^*)] \leq 0,$$

where the infinite sum exists by Assumption A.5(iv). By Assumption A.5(iii), for (A.14) to hold, we must have

$$(A.15) \quad \sum_{t=1}^{\infty} [r_t(y_t^* + \delta, y_{t+1}^* + \delta) - r_t(y_t^*, y_{t+1}^*)] < \infty.$$

Dividing through by δ and recalling (A.3), we get $\sum_{t=1}^{\infty} m_t(-\delta) < \infty$. Thus by (A.13) and Theorem A.1(ii), $\overline{\lim}_{t \uparrow \infty} q_t \leq 0$. \square

A.2 Proof of Theorem 2.1

Let $\{x_t\}$ be a feasible path satisfying (2.7) and (2.8). Then

$$(A.16) \quad \overline{\lim}_{T \uparrow \infty} \sum_{t=0}^T [V_t(x_t^*, x_{t+1}^*) - V_t(x_t, x_{t+1})] < \infty.$$

It follows that

$$(A.17) \quad \{V_t(x_t^*, x_{t+1}^*) - V_t(x_t, x_{t+1})\}_{t=1}^{\infty} \in \Psi,$$

where Ψ is defined by (A.8). For $t \in \mathbb{Z}_+$, let $e_t = x_t^* - x_t$. Consider the following problem.

$$(A.18) \quad \left\{ \begin{array}{l} \max_{\{y_t\}_{t=0}^{\infty}} \sum_{t=0}^{\infty} V_t(x_t^* + y_t e_t, x_{t+1}^* + y_{t+1} e_{t+1}) \\ \text{s.t. } y_0 = 0, \quad \forall t \in \mathbb{Z}_+, (x_t^* + y_t e_t, x_{t+1}^* + y_{t+1} e_{t+1}) \in X_t, \end{array} \right.$$

where $\forall t \in \mathbb{Z}_+$, $y_t \in \mathbb{R}$. For $t \in \mathbb{Z}_+$, define

$$(A.19) \quad r_t(y_t, y_{t+1}) = V_t(x_t^* + y_t e_t, x_{t+1}^* + y_{t+1} e_{t+1}),$$

$$(A.20) \quad Y_t = \{(y_t, y_{t+1}) \in \mathbb{R}^2 \mid (x_t^* + y_t e_t, x_{t+1}^* + y_{t+1} e_{t+1}) \in X_t\}.$$

Obviously Assumptions A.1 and A.2 hold. For $t \in \mathbb{Z}_+$, let $y_t^* = 0$. Then $\{y_t^*\}$ is optimal for (A.18) since $\{x_t^*\}$ is optimal for (2.1). Thus Assumption A.3 holds. Assumption A.4 follows from the convexity of X_t and (2.8). Note that $\forall t \in \mathbb{Z}_+$,

$$(A.21) \quad q_t = \lim_{\epsilon \downarrow 0} \frac{V_t(x_t^*, x_{t+1}^* - \epsilon e_t) - V_t(x_t^*, x_{t+1}^*)}{\epsilon} = V_{t,2}(x_t^*, x_{t+1}^*; -e_t).$$

Note also that $\forall t \in \mathbb{Z}_+, \forall \epsilon \in (0, 1]$,

$$(A.22) \quad m_t(\epsilon) = \frac{V_t(x_t^*, x_{t+1}^*) - V_t(x_t^* - \epsilon e_t, x_{t+1}^* - \epsilon e_{t+1})}{\epsilon}$$

$$(A.23) \quad \leq V_t(x_t^*, x_{t+1}^*) - V_t(x_t^* - e_t, x_{t+1}^* - e_{t+1}),$$

where the inequality holds by concavity. Recalling (A.17) and the definition of $\{e_t\}$, we see that (A.12) holds with $b_t = V_t(x_t^*, x_{t+1}^*) - V_t(x_t, x_{t+1})$. Thus TVC (2.6) holds by Theorem A.1(i) and (A.21).

A.3 Proof of Theorem 2.2

Consider the following problem.

$$(A.24) \quad \begin{cases} \max_{\{y_t\}_{t=0}^{\infty}} \sum_{t=0}^{\infty} V_t(y_t x_t^*, y_{t+1} x_{t+1}^*) \\ \text{s.t. } y_0 = 1, \quad \forall t \in \mathbb{Z}_+, (y_t x_t^*, y_{t+1} x_{t+1}^*) \in X_t, \end{cases}$$

where $\forall t \in \mathbb{Z}_+$, $y_t \in \mathbb{R}$. For $t \in \mathbb{Z}_+$, define

$$(A.25) \quad r_t(y_t, y_{t+1}) = V_t(y_t x_t^*, y_{t+1} x_{t+1}^*),$$

$$(A.26) \quad Y_t = \{(y_t, y_{t+1}) \in \mathbb{R}^2 \mid (y_t x_t^*, y_{t+1} x_{t+1}^*) \in X_t\}.$$

Obviously Assumptions A.1 and A.2 hold. For $t \in \mathbb{Z}_+$, let $y_t^* = 1$. Then $\{y_t^*\}$ is optimal for (A.24) since $\{x_t^*\}$ is optimal for (2.1). Thus Assumption A.3 holds. Assumption A.4 follows from Assumption 2.4. Note that $\forall t \in \mathbb{Z}_+$,

$$(A.27) \quad q_t = \lim_{\epsilon \downarrow 0} \frac{V_t(x_t^*, (1 - \epsilon)x_{t+1}^*) - V_t(x_t^*, x_{t+1}^*)}{\epsilon} = V_{t,2}(x_t^*, x_{t+1}^*; -x_{t+1}^*).$$

Note also that $\forall t \in \mathbb{N}, \hat{m}_t(1 - \lambda) = \hat{w}_t(\lambda)$. Thus (A.12) holds by (2.12). Hence both conclusions hold by Theorem A.1.

A.4 Proof of Theorem 2.3

Consider (A.24) again. For $t \in \mathbb{Z}_+$, define r_t and Y_t by (A.25) and (A.26). It is easy to see that Assumptions A.1–A.5 hold. Note that $\forall t \in \mathbb{N}, \hat{m}_t(1 - \lambda) = \hat{w}_t(\lambda)$ and $m_t(1 - \mu) = w_t(\mu)$. Thus (A.13) holds by (2.18). Recalling (A.27), we see that TVC (2.16) holds by Theorem A.2.

A.5 Proof of Theorem 3.1

By Theorem 2.1, TVC (2.6) holds. Thus it suffices to verify (3.8). Let $t \in \mathbb{Z}_+$ and $e_t = x_t^* - x_t$. For $\tilde{\epsilon} \in (0, \epsilon]$ (ϵ is given by (3.7)), define

$$(A.28) \quad a_t(\tilde{\epsilon}) = \frac{v_t(x_t^*, x_{t+1}^* - \tilde{\epsilon}e_{t+1}) - v_t(x_t^*, x_{t+1}^*)}{\tilde{\epsilon}}.$$

By (3.7), $Ea_t(\epsilon) > -\infty$. By concavity, $a_t(\tilde{\epsilon})$ is nonincreasing in $\tilde{\epsilon} \in (0, \epsilon]$. Hence $\forall \tilde{\epsilon} \in (0, \epsilon], a_t(\tilde{\epsilon}) \geq a_t(\epsilon)$. Now by the monotone convergence theorem, $\lim_{\tilde{\epsilon} \downarrow 0} Ea_t(\tilde{\epsilon}) = E \lim_{\tilde{\epsilon} \downarrow 0} a_t(\tilde{\epsilon})$, which is equivalent to (3.8).

A.6 Proof of Theorem 3.2

Immediate from Theorem 2.2 and Assumption 3.3.

A.7 Proof of Theorem 3.3

Immediate from Theorem 2.3 and Assumption 3.3.

A.8 Proof of Proposition 4.1

Note from Remark 3.3 that

$$(A.29) \quad \text{Assumption 4.3} \Rightarrow \text{Assumption 3.3.}$$

Thus to conclude TVC (4.1) from Theorem 3.2 and Remark 4.1, it suffices to verify (2.12) and (2.15). Let $\underline{\lambda} \in (\underline{\lambda}_0, 1)$, where $\underline{\lambda}_0$ is given by Assumption 2.4. Let $t \in \mathbb{N}$. By Assumptions

4.1 and 4.2,

$$(A.30) \quad 0 \leq \frac{\partial Ev_t(\lambda x_t^*, \lambda x_{t+1}^*)}{\partial \lambda} \Big|_{\lambda=1} = \frac{\partial \lambda^\alpha Ev_t^*}{\partial \lambda} \Big|_{\lambda=1} = \alpha Ev_t^*,$$

where $v_t^* = v_t(x_t^*, x_{t+1}^*)$. By Assumption 4.1 and (A.30),

$$(A.31) \quad \forall \lambda \in [\underline{\lambda}, 1], \quad w_t(\lambda) = \frac{1 - \lambda^\alpha}{(1 - \lambda)\alpha} \alpha Ev_t^* \leq A\alpha Ev_t^*,$$

where $A = \sup_{\lambda \in [\underline{\lambda}, 1]} (1 - \lambda^\alpha)/[(1 - \lambda)\alpha] \in (0, \infty)$; A is finite since $(1 - \lambda^\alpha)/[(1 - \lambda)\alpha]$ is continuous on $[\underline{\lambda}, 1]$ by l'Hôpital's rule. Now (2.12) and (2.15) follow from (A.31) and (4.2).

A.9 Proof of Proposition 4.2

Recall (A.29). To conclude TVC (4.1) from Theorem 3.3 and Remark 4.1, it suffices to verify (2.18). Let $\mu \in (1, \bar{\mu}]$, where $\bar{\mu}$ is given by Assumption 2.5. By Assumption 4.1,

$$(A.32) \quad \forall t \in \mathbb{N}, \quad w_t(\mu) = \frac{\mu^\alpha - 1}{(\mu - 1)\alpha} \alpha Ev_t^*.$$

Now (2.18) follows from (A.31) and (A.32).

A.10 Proof of Proposition 4.3

Note from Remarks 2.3 and 3.3 that

$$(A.33) \quad \text{Assumptions 4.5 and 4.6} \Rightarrow \text{Assumption 2.4 and 3.3.}$$

To conclude TVC (4.1) from Therem 3.2 and Remark 4.2, it suffices to verify (2.12) and (2.15). Let $\lambda \in (0, 1)$, $\lambda \in [\underline{\lambda}, 1)$, and $t \in \mathbb{N}$. By concavity and Assumption 4.5(iii),

$$(A.34) \quad g_t(\lambda x_t^*, \lambda x_{t+1}^*) \geq \lambda g_t(x_t^*, x_{t+1}^*) + (1 - \lambda)g_t(0, 0) \geq \lambda g_t(x_t^*, x_{t+1}^*).$$

We have

$$(A.35) \quad (1 - \lambda)\beta^{-t}w_t(\lambda) = E[\ln g_t(x_t^*, x_{t+1}^*) - \ln g_t(\lambda x_t^*, \lambda x_{t+1}^*)]$$

$$(A.36) \quad \leq E[\ln g_t(x_t^*, x_{t+1}^*) - \ln(\lambda g_t(x_t^*, x_{t+1}^*))] = -\ln \lambda,$$

where the inequality holds by (A.34). It follows that

$$(A.37) \quad w_t(\lambda) \leq \beta^t \frac{-\ln \lambda}{1 - \lambda} \leq \beta^t \frac{-\ln \underline{\lambda}}{1 - \underline{\lambda}}.$$

Now (2.12) and (2.15) follow.

A.11 Proof of Proposition 4.4

Recall (A.33). To conclude TVC (4.1) from Therem 3.2 and Remark 4.2, it suffices to verify (2.12) and (2.15). Let $\underline{\lambda} \in (0, 1)$, $\lambda \in [\underline{\lambda}, 1]$, and $t \in \mathbb{N}$. We have

$$(A.38) \quad (1 - \lambda)\beta^{-t}w_t(\lambda) = E[u(g_t(x_t^*, x_{t+1}^*)) - u(g_t(\lambda x_t^*, \lambda x_{t+1}^*))]$$

$$(A.39) \quad \leq E[u(g_t(x_t^*, x_{t+1}^*)) - u(\lambda g_t(x_t^*, x_{t+1}^*))]$$

$$(A.40) \quad = (1 - \lambda^\alpha)Eu(g_t(x_t^*, x_{t+1}^*)),$$

where (A.39) holds by (A.34), and (A.40) holds by Assumption 4.4. It follows that

$$(A.41) \quad w_t(\lambda) \leq \beta^t \frac{1 - \lambda^\alpha}{(1 - \lambda)\alpha} \alpha Eu(g_t^*) \leq \beta^t A\alpha Eu(g_t^*),$$

where $g_t^* = g_t(x_t^*, x_{t+1}^*)$ and A is as in (A.31). Now (2.12) and (2.15) follow from (4.6) and (A.41).²²

B A Further Result on the Model of Section 4.2

This appendix considers the model of Section 4.2 and shows a result that does not assume (4.6), i.e., the finiteness of the objective function at the optimum.

Proposition B.1. *Assume Assumptions 2.3, 2.5, 3.1, 3.2, and 4.4–4.6. Suppose $\alpha \neq 0$. For $\lambda \in [0, \bar{\mu}]$, let $\gamma_t(\lambda) = g_t(\lambda x_t^*, \lambda x_{t+1}^*)$, where $\bar{\mu}$ is given by Assumption 2.5. Let $\bar{\epsilon} = \min\{1, \bar{\mu} - 1\}$. Suppose*

$$(B.1) \quad \exists \epsilon \in (0, \bar{\epsilon}), \exists \theta \geq 0, \forall t \in \mathbb{N}, \quad 0 \leq \gamma_t(1) - \gamma_t(1 - \epsilon) \leq \theta[\gamma_t(1 + \epsilon) - \gamma_t(1)].$$

Then TVC (4.5) holds.²³

²²This proof works without Assumption 4.4 if u is nondecreasing, concave, and differentiable and if $\exists \lambda \in (0, 1), \exists \alpha \in (-\infty, 1], \forall t \in \mathbb{N}, \forall \lambda \in [\underline{\lambda}, 1], u(\lambda g_t^*) \geq \lambda^\alpha u(g_t^*)$. In this case, “=” in (A.40) must be replaced by “≤.” That $\alpha Eu(g_t^*) \geq 0$ can be shown from the inequality $u(\lambda g_t^*) \geq \lambda^\alpha u(g_t^*)$. Note that both sides are equal at $\lambda = 1$. Thus, differentiating both sides with respect to λ and evaluating at $\lambda = 1$, we have $0 \leq u'(g_t^*)g_t^* \leq \alpha u(g_t^*)$.

²³Proposition B.1 can easily be generalized to more general utility functions. In fact the proof of Proposition B.1 works without Assumption 4.4 if u is nondecreasing, concave, and differentiable and if $\exists \sigma \geq 0, \forall t \in \mathbb{N}, u'(\gamma_t(1 - \epsilon)) \leq \sigma u'(\gamma_t(1 + \epsilon))$, where ϵ is given by (B.1). (To verify this inequality, one may utilize (A.34) and (B.3).) In this case, the first inequalities in (B.2) and (B.4) together with (B.1) imply (2.18).

Proof. Recall (A.33). To conclude TVC (4.1) from Theorem 3.3 and Remark 4.2, it suffices to verify (2.18). Let $\lambda = 1 - \epsilon$, $\mu = 1 + \epsilon$, and $t \in \mathbb{N}$. Note from (A.38) that by concavity,

$$(B.2) \quad w_t(\lambda) \leq \frac{\beta^t Eu'(\gamma_t(\lambda))[\gamma_t(1) - \gamma_t(\lambda)]}{1 - \lambda} \leq \frac{\beta^t \lambda^{\alpha-1} Eu'(\gamma_t(1))[\gamma_t(1) - \gamma_t(\lambda)]}{1 - \lambda},$$

where the second inequality uses (A.34), the first inequality in (B.1), and Assumption 4.4. By concavity, $\gamma_t(1) \geq (1 - (1/\mu))\gamma_t(0) + \gamma_t(\mu)/\mu \geq \gamma_t(\mu)/\mu$ (the second inequality holds by Assumption 4.5(iii)); thus

$$(B.3) \quad \gamma_t(\mu) \leq \mu\gamma_t(1).$$

By concavity,

$$(B.4) \quad w_t(\mu) \geq \frac{\beta^t Eu'(\gamma_t(\mu))[\gamma_t(\mu) - \gamma_t(1)]}{\mu - 1} \geq \frac{\beta^t \mu^{\alpha-1} Eu'(\gamma_t(1))[\gamma_t(\mu) - \gamma_t(1)]}{\mu - 1},$$

where the second inequality uses (B.3) and Assumption 4.4. Recalling Remark 2.4, we obtain (2.18) from (B.1), (B.2), and (B.4). \square

Note that (B.1) holds with $\theta = 1$ if $\exists n \in \mathbb{N}, \forall t \in \mathbb{N}$, (i) $B_t = \mathbb{R}^n$ (recall Assumption 3.1), (ii) $g_t(x_t^*, x_{t+1}^*) \geq g_t(0, 0)$, and (iii) $\forall (y, z) \in X_t, g_t(y, z) = a_t + b_t y + c_t z$ for some $a_t : \Omega \rightarrow \mathbb{R}$ and $b_t, c_t : \Omega \rightarrow \mathbb{R}^n$. These conditions are satisfied in single-agent asset pricing models of the type studied by Lucas (1978), Kamihigashi (1998), and Montrucchio and Privileggi (2000). For such models, TVC (4.5) can be used to rule out bubbles.²⁴ Even if g_t is nonlinear, condition (B.1) can be satisfied; e.g., it is satisfied in a stochastic discrete-time version of the asset pricing model with nonlinear constraints discussed in Kamihigashi (2000b).

References

- Alvarez, F., and N.L. Stokey, 1998, “Dynamic Programming with Homogeneous Functions,” *Journal of Economic Theory* 82, 167–189.
- Brock, W.A., 1970, “On Existence of Weakly Maximal Programmes in a Multi-Sector Economy,” *Review of Economic Studies* 37, 275–280.
- Brock, W.A., and L.J. Mirman, 1972, “Optimal Economic Growth and Uncertainty: The Discounted Case,” *Journal of Economic Theory* 4, 479–513.

²⁴The TVC by itself does not rule out bubbles since the Euler equation may be violated; see Kamihigashi (1998) and Montrucchio and Privileggi (2000). See Santos and Woodford (1997) and Montrucchio (2000) for results on bubbles in economies with heterogeneous agents.

- Ekeland, I., and J.A. Scheinkman, 1986, “Transversality Conditions for Some Infinite Horizon Discrete Time Optimization Problems,” *Mathematics of Operations Research* 11, 216–229.
- Halkin, H., 1974, “Necessary Conditions for Optimal Control Problems with Infinite Horizons,” *Econometrica* 42, 267–272.
- Kamihigashi, T., 1998, “Uniqueness of Asset Prices in an Exchange Economy with Unbounded Utility,” *Economic Theory* 12, 103–122.
- Kamihigashi, T., 2000a, “A Simple Proof of Ekeland and Scheinkman’s Result on the Necessity of a Transversality Condition,” *Economic Theory* 15, 463–468.
- Kamihigashi, T., 2000b, “Necessity of Transversality Conditions for Infinite Horizon Problems,” *Econometrica*, forthcoming.
 (Downloadable at http://papers.ssrn.com/paper.taf?ABSTRACT_ID=233633)
- King, R.G., C.I. Plosser, and S.T. Rebelo, 1988, “Production, Growth and Business cycles: I. The basic Neoclassical Model,” *Journal of Monetary Economics* 21, 195–232.
- Lucas, R.E., Jr., 1978, “Asset Prices in an Exchange Economy,” *Econometrica* 46, 1429–1445.
- Lucas, R.E., Jr., 1988, “On the Mechanics of Economic Development,” *Journal of Monetary Economics* 22, 3–42.
- Michel, P., 1990, “Some Clarifications on the Transversality Condition,” *Econometrica* 58, 705–723.
- Montrucchio, L., 2000, “Optimality Necessary Conditions and Bubbles in Sequential Economies with Heterogeneous Agents,” University of Turin.
- Montrucchio, L., and F. Privileggi, 2000, “On Fragility of Bubbles in Equilibrium Asset Pricing Models of Lucas-Type,” *Journal of Economic Theory*, forthcoming.
- Peleg, B., 1970, “Efficiency Prices for Optimal Consumption Plans III,” *Journal of Mathematical Analysis and Applications* 32, 630–638.
- Peleg, B., and H.E. Ryder, Jr., 1972, “On Optimal Consumption Plans in a Multi-sector Economy,” *Review of Economic Studies* 39, 159–169.
- Rebelo, S. (1991), “Long-Run Policy Analysis and Long-Run Growth,” *Journal of Political Economy* 99, 500–521.
- Santos, M., and M. Woodford, 1997, “Rational Asset Pricing Bubbles,” *Econometrica* 65, 19–57.
- Shell, K., 1969, “Applications of Pontryagin’s Maximum Principle to Economics,” in *Mathematical Systems Theory and Economics I*, edited by H.W. Kuhn and G.P. Szegö. New York: Springer-Verlag, pp. 241–292.
- Takekuma, S., 1992, “Optimal Growth under Uncertainty: A Complete Characterization of Weakly Maximal Programs,” *Hitotsubashi Journal of Economics* 33, 169–182.
- Weitzman, M.L., 1973, “Duality Theory for Infinite Horizon Convex Models,” *Management Science* 19, 783–789.

- Yano, M., 1989, “Comparative Statics in Dynamic Stochastic Models: Differential Analysis of a Stochastic Modified Golden Rule State in a Banach Space,” *Journal of Mathematical Economics* 18, 169–185.
- Zilcha, I., 1976, “Characterization by Prices of Optimal Programs under Uncertainty,” *Journal of Mathematical Economics* 3, 173–181.
- Zilcha, I., 1978, “Transversality Condition in a Multi-sector Economy under Uncertainty,” *Econometrica* 46, 515–525.